



**NATIONAL HEALTH  
LABORATORY SERVICE**

NATIONAL INSTITUTE FOR OCCUPATIONAL HEALTH

---

## **Pathology Division Surveillance Report**

### **Demographic Data and Disease Rates for January to December 1995**

**Ntombizodwa Ndlovu**

**Tony Davies**

**Gill Nelson**

**Jill Murray**

*PO Box 4788  
Johannesburg  
2000*

*e-mail: [jill.murray@nioh.nhls.ac.za](mailto:jill.murray@nioh.nhls.ac.za)*

**NIOH Report 28/2011**

[http://www.nioh.ac.za/publications/publications\\_pathaut\\_reports.htm](http://www.nioh.ac.za/publications/publications_pathaut_reports.htm)

## **ACKNOWLEDGEMENTS**

*We thank all the staff members of the NIOH Pathology Department, for their invaluable contribution to the autopsy service.*

*We also thank Prof Tony Davies for editing the manuscript.*

*National Institute for Occupational Health, PO Box 4788, Johannesburg, 2000, South Africa*


*ISSN 1812-7681*

## EXECUTIVE SUMMARY

---

During 1995, 4 003 cases came to autopsy at the NIOH. Of these, 70.7% were black men, 26.5% were white, 2.5% were coloured and for 12 cases the population group was not known.

Overall disease rates (per 1000 autopsies) for 1995 are shown in Figure 1.



**FIG 1 OVERALL DISEASE RATES FOR 1995**

# TABLE OF CONTENTS

<b>EXECUTIVE SUMMARY</b>	<i>i</i>
<b>GLOSSARY</b>	<i>iv</i>
<b>SECTION 1: BACKGROUND</b>	1
<b>SECTION 2: DEMOGRAPHIC DATA</b>	2
<b>SECTION 3: ACTIVE TUBERCULOSIS</b>	7
<b>SECTION 4: SILICOSIS</b>	9
<b>SECTION 5: OTHER PNEUMOCONIOSES</b>	
<i>Massive Fibrosis</i>	12
<i>Coal Workers' Pneumoconiosis</i>	12
<i>Mixed Dust Pneumoconiosis</i>	12
<i>Asbestosis and pleural plaques</i>	12
<b>SECTION 6: EMPHYSEMA</b>	13
<b>SECTION 7: MESOTHELIOMA</b>	15
<b>SECTION 8: PRIMARY LUNG CANCER</b>	16
<b>SECTION 9: CLINICAL CAUSES OF DEATH</b>	17

## LIST OF TABLES

<b>Table 2.1</b>	<i>Distribution of autopsies by year and population group (1995)</i>	2
<b>Table 2.2</b>	<i>Number and proportion of autopsies by type and population group (1995)</i>	3
<b>Table 2.3</b>	<i>Number and proportion of autopsies by age and population group (1995)</i>	3
<b>Table 2.4</b>	<i>Number and proportion of autopsies by commodity and population group (1995)</i>	4
<b>Table 2.5</b>	<i>Number and proportion of autopsies by years of service and population group (1995)</i>	5
<b>Table 2.6</b>	<i>Mean age by commodity and population group (1995)</i>	6
<b>Table 2.7</b>	<i>Mean duration of service by commodity and population group (1995)</i>	7
<b>Table 3.1</b>	<i>Number of cases and prevalence of active PTB by commodity and population group (1995)</i>	9
<b>Table 3.2</b>	<i>Number of cases and prevalence of active PTB by age and population group (1995)</i>	9
<b>Table 4.1</b>	<i>Number of cases and prevalence of silicosis by commodity and population group (1995)</i>	10
<b>Table 4.2</b>	<i>Number of cases and prevalence of silicosis in the gold mining industry, by age and population group (1995)</i>	10
<b>Table 4.3</b>	<i>Number of cases and prevalence of silicosis in the gold mining industry, by years of service and population group (1995)</i>	11
<b>Table 5.1</b>	<i>Number of cases and prevalence of asbestosis by age and population group (1995)</i>	12

<b>Table 6.1</b>	<i>Number of cases and prevalence of emphysema by age and population group (1995)</i>	13
<b>Table 6.2</b>	<i>Number of cases and prevalence of emphysema by commodity and population group (1995)</i>	13
<b>Table 6.3</b>	<i>Number of cases and prevalence of emphysema by years of service and population group (1995)</i>	14
<b>Table 7.1</b>	<i>Number and proportion of mesothelioma cases by age and population group (1995)</i>	15
<b>Table 7.2</b>	<i>Number and proportion of mesothelioma cases by commodity and population group (1995)</i>	15
<b>Table 8.1</b>	<i>Number of cases and prevalence of primary lung cancer by age and population group (1995)</i>	16
<b>Table 8.2</b>	<i>Number of cases and prevalence of primary lung cancer by commodity and population group (1995)</i>	16
<b>Table 9.1</b>	<i>Clinical causes of death by population group (1995)</i>	17

## LIST OF FIGURES

<b>Figure 1</b>	<i>Overall disease rates for 1995</i>	<i>i</i>
<b>Figure 2.1</b>	<i>Distribution of autopsies by age and population group (1995)</i>	4
<b>Figure 2.2</b>	<i>Distribution of autopsies by commodity and population group (1995)</i>	5
<b>Figure 2.3</b>	<i>Distribution of autopsies by years of service and population group (1995)</i>	6
<b>Figure 3.1</b>	<i>Distribution of active TB by site (1995)</i>	8
<b>Figure 9.1</b>	<i>Clinical cause of death as given by the clinicians who submit the organs to the NIOH (1995)</i>	17

## GLOSSARY

---

<b>Asbestosis</b>	<i>Lung fibrosis caused by asbestos fibres</i>
<b>Coal workers' pneumoconiosis</b>	<i>Lung fibrosis caused by exposure to coal dust</i>
<b>Emphysema</b>	<i>Lung disease caused by the destruction of the alveolar walls</i>
<b>Massive fibrosis</b>	<i>Lung fibrosis caused by exposure to dust and measuring more than 1 cm in diameter</i>
<b>Mesothelioma</b>	<i>A malignant tumour of the pleural cavity of the lungs</i>
<b>Miner</b>	<i>A person who has worked in a controlled mine or works</i>
<b>Mixed exposures</b>	<i>The multiple dust types to which a miner may be exposed, having worked in several mining commodities in his lifetime</i>
<b>Prevalence</b>	<i>The number of cases in a defined population at a given time</i>
<b>Silicosis</b>	<i>Lung fibrosis caused by inhalation of silica dust; detected by the presence of silicotic nodules in the lung tissue</i>
<b>Surveillance</b>	<i>The ongoing and systematic collection, analysis and interpretation of data related to adverse health outcomes</i>

## SECTION 1 – BACKGROUND

---

*The Occupational Diseases in Mines and Works Act, 1973 (Act 78 of 1973) requires that the cardio-respiratory organs of a deceased person who has worked at a controlled mine or a controlled works be examined for the presence of occupational disease, regardless of the clinical cause of death and provided that the next of kin agrees. These examinations are performed by pathologists at the National Institute for Occupational Health (NIOH). A detailed report on each case examined is sent to the Medical Bureau for Occupational Diseases (MBOD). Cases certified as having a compensable disease are then referred to the Compensation Commissioner's office, where the payment for compensation is managed.*

*Since 1975, the pathological findings from the autopsy examinations have been recorded on the computerised PATHAUT database. PATHAUT comprises data from autopsy examinations and clinical files which include occupational histories. The database is unique and provides an important resource for both surveillance and research. These data are the only comprehensive surveillance data on occupational lung disease in the South African mining industry.*

*The data presented in this report summarise the PATHAUT system surveillance results, i.e. the results of the systematic collection, collation, and analysis of the pathology findings in the cardio-respiratory organs of mine workers. Data from PATHAUT are exported into, and analysed, using SAS v9.1.*

*This report describes autopsy cases examined during the year 1995. This and other annual reports can be accessed at [www.nioh.ac.za](http://www.nioh.ac.za).*

*Many of the cases had "mixed" exposures in that they had been employed in more than one commodity. For simplicity, cases are categorised according to the commodity in which most years of service were recorded, i.e. the commodity in which the miners had worked for the longest period.*

*All disease rates reported in this document are expressed per 1000. In all calculations, the denominators used are the total numbers of autopsies in specific commodities, age groups or population groups. Some of these rates must be viewed with caution, as the denominators are very small. This applies, for example, to those commodities where few workers are employed (such as manganese mining), and to the older age groups in some instances.*

## SECTION 2 – DEMOGRAPHIC DATA

The number of autopsies performed for 1995 is presented in Table 2-1.

**TABLE 2-1 DISTRIBUTION OF AUTOPSIES BY YEAR AND POPULATION GROUP (1975-1995)**

Year of autopsy	Black		White		Coloured		Unknown		Total N
	N	%	N	%	N	%	N	%	
1975	2 190	71	854	28	32	1			3 076
1976	2 335	68	1 072	31	27	1			3 434
1977	2 351	69	1 039	30	33	1			3 423
1978	2 245	67	1 090	32	32	1			3 367
1979	2 118	66	1 026	33	45	1			3 189
1980	2 338	64	1 274	35	46	1			3 658
1981	2 209	66	1 117	33	33	1			3 359
1982	2 312	63	1 302	36	44	1	1		3 659
1983	2 096	65	1 109	34	41	1			3 246
1984	1 966	64	1 098	36	28	1			3 092
1985	2 275	64	1 200	34	66	2			3 541
1986	2 456	68	1 125	31	45	1			3 626
1987	2 594	68	1 168	30	78	2			3 840
1988	2 518	67	1 165	31	77	2			3 760
1989	2 138	65	1 090	33	60	2			3 288
1990	2 172	64	1 155	34	51	2			3 378
1991	2 143	65	1 080	33	66	2			3 289
1992	2 144	66	1 049	32	70	2			3 263
1993	1 863	65	956	33	65	2			2 884
1994	1 737	61	1 021	36	94	3			2 852
1995	2 830	71	1 062	27	99	2	12	0.3	4 003
<b>Total</b>	<b>47 030</b>	<b>66</b>	<b>23 052</b>	<b>32</b>	<b>1 132</b>	<b>2</b>	<b>13</b>		<b>71 227</b>

*It is important to note that a referral bias exists: there is a low autopsy rate for black men who have left employment at the mines, whereas the majority of white retired miners come to autopsy.*

*The pathologists at the NIOH perform two types of autopsy examinations. For men dying distant from Johannesburg, the cardio-respiratory organs are removed locally, preserved in formalin and sent to the NIOH. Full autopsies are undertaken on men who die close to Johannesburg.*



Table 2-2 shows the distribution of autopsies by population group for 1995. Autopsies of only the cardio-respiratory organs comprised 91.5% of all examinations.


**TABLE 2-2 NUMBER AND PROPORTION OF AUTOPSIES BY TYPE AND POPULATION GROUP (1995)**

Autopsy type	Black		White		Coloured		Unknown		Total	
	N	%	N	%	N	%	N	%	N	%
Cardio-respiratory organs only	2 755	97.3	799	75.2	98	99.0	12	100	3 664	91.5
Full autopsy	75	2.7	262	24.7	0	-	0	-	337	8.4
Not stated	0	-	1	0.1	1	1.0	0	-	2	0.1
<b>Total</b>	2 830		1 062		99		12		4 003	

The age distributions of autopsies for 1995 are shown in Table 2-3 and Figure 2-1. The mean age at autopsy of black men was 36.1 years. The mean age of white men at autopsy was 60.1 years and for coloured men 62.7 years

**TABLE 2-3 NUMBER AND PROPORTION OF AUTOPSIES BY AGE AND POPULATION GROUP (1995)**

Age group (years)	Black		White		Coloured		Unknown		Total	
	N	%	N	%	N	%	N	%	N	%
<20	3	0.1	2	0.2	0	-	0	-	5	0.1
20-29	278	9.8	38	3.6	2	2.0	0	-	318	7.9
30-39	898	31.7	80	7.5	5	5.1	2	16.7	985	24.6
40-49	775	27.4	115	10.8	10	10.1	1	8.3	901	22.5
50-59	463	16.4	192	18.1	18	18.2	0	-	673	16.8
60-69	58	2.0	267	25.1	31	31.3	0	-	356	8.9
70-79	7	0.2	252	23.7	17	17.2	0	-	276	6.9
80+	6	0.2	100	9.4	15	15.2	0	-	121	3.0
Unknown	342	12.1	16	1.5	1	1.0	9	75.0	368	9.2
<b>Total</b>	2 830		1 062		99		12		4 003	



**FIG 2-1 DISTRIBUTION OF AUTOPSIES BY AGE AND POPULATION GROUP (1995)**


Cases were placed in categories according to the commodity in which they had worked for the longest duration (most exposure). Many men worked in a number of different mining commodities during their lifetimes and had “mixed” exposures. This was not taken into account in the analysis of exposure type (commodity).

Table 2-4 and Figure 2-2 show the distribution of autopsies by commodity and population group for 1995.

**TABLE 2-4 NUMBER AND PROPORTION OF AUTOPSIES BY COMMODITY AND POPULATION GROUP (1995)**

Commodity	Black		White		Coloured		Unknown		Total	
	N	%	N	%	N	%	N	%	N	%
Gold	2 313	81.7	771	72.6	4	4.0	2	16.7	3 090	77.2
Platinum	251	8.9	44	4.1	0	-	1	8.3	296	7.4
Coal	121	4.3	73	6.9	1	1.0	0	-	195	4.9
Asbestos	30	1.1	26	2.4	74	74.7	0	-	130	3.2
Iscor	12	0.4	50	4.7	1	1.0	0	-	63	1.6
Diamond	11	0.4	20	1.9	1	1.0	0	-	32	0.8
Copper	1	-	27	2.5	15	15.2	0	-	43	1.1
Other	5	0.2	2	0.2	0	-	0	-	7	0.2
Unknown	86	3.0	49	4.6	3	3.0	9	75.0	147	3.7
<b>Total</b>	<b>2 830</b>		<b>1 062</b>		<b>99</b>		<b>12</b>		<b>4 003</b>	

NOTE: This table shows only those commodities where a total of 10 or more cases were received




**FIG 2-2 DISTRIBUTION OF AUTOPSIES BY COMMODITY AND POPULATION GROUP (1995)**

Detailed information about the years in mining service by population group is presented in Table 2-5 and Figure 2-3.

**TABLE 2-5 NUMBER AND PROPORTION OF AUTOPSIES BY YEARS OF SERVICE AND POPULATION GROUP (1995)**

Years of service	Black		White		Coloured		Unknown		Total	
	N	%	N	%	N	%	N	%	N	%
<1	40	1.4	2	0.2	0	-	0	-	42	1.0
1-5	290	10.2	101	9.5	15	15.2	0	-	406	10.1
6-10	452	16.0	100	9.4	23	23.2	1	8.3	576	14.4
11-15	466	16.5	123	11.6	17	17.2	1	8.3	607	15.2
16-20	366	12.9	122	11.5	14	14.1	1	8.3	503	12.6
21-25	98	3.5	132	12.4	6	6.1	0	-	236	5.9
26-30	52	1.8	117	11.0	5	5.1	0	-	174	4.3
31-35	18	0.6	103	9.7	0	-	0	-	121	3.0
36-40	12	0.4	63	5.9	3	3.0	0	-	78	1.9
41+	0	-	26	2.4	1	1.0	0	-	27	0.7
Unknown	1 036	36.6	173	16.3	15	15.2	9	75.0	1 233	30.8
<b>Total</b>	<b>2 830</b>		<b>1 062</b>		<b>99</b>		<b>12</b>		<b>4 003</b>	



**FIG 2-3 DISTRIBUTION OF AUTOPSIES BY YEARS OF SERVICE AND POPULATION GROUP (1995)**

The mean age and duration of service by commodity type and population group for those cases for which information was available are shown in Tables 2-6 and 2-7.

**TABLE 2-6 MEAN AGE BY COMMODITY AND POPULATION GROUP (1995)**

Commodity	Black			White		
	N	Mean (years)	SD*	N	Mean (years)	SD*
Gold	2 032	40.8	9.7	764	62.2	15.2
Platinum	236	41.6	9.2	44	58.3	15.9
Coal	110	41.0	9.9	73	54.8	17.0
Asbestos	29	54.0	11.3	26	62.3	11.7
Iscor	12	43.7	8.3	50	55.8	14.8
Diamond	9	36.0	10.3	19	56.4	11.8
Copper	1	63	-	27	62.9	13.6
Other	5	54.6	5.8	2	72.0	15.6
Unknown	54	40.9	10.7	41	59.4	16.7
Total	2 488	41.1	9.8	1 046	61.1	15.4

\* Standard deviation


**TABLE 2-7 MEAN DURATION OF SERVICE BY COMMODITY AND POPULATION GROUP (1995)**

Commodity	Black			White		
	N	Mean (years)	SD*	N	Mean (years)	SD*
<i>Gold</i>	1 659	12.6	7.0	674	22.2	11.3
<i>Platinum</i>	70	9.0	5.3	34	15.4	8.1
<i>Coal</i>	33	12.3	7.6	61	15.5	10.9
<i>Asbestos</i>	15	12.8	8.8	21	12.6	9.6
<i>Iscor</i>	3	17.1	3.0	39	17.6	9.6
<i>Diamond</i>	7	9.5	10.8	17	21.9	20.5
<i>Copper</i>	0	-	-	25	16.9	8.8
<i>Other</i>	4	18.7	9.3	2	13.4	2.0
<i>Unknown</i>	3	21.7	15.2	16	17.2	10.4
<i>Total</i>	1 794	12.5	7.0	889	20.8	11.5

\* Standard deviation

### SECTION 3 – ACTIVE TUBERCULOSIS

The distribution of active tuberculosis (TB) by anatomical site is presented in Figure 3-1 (n=458). Active pulmonary TB (PTB) was diagnosed in 9.3% (n=373) of all cases autopsied in 1995. Most of the men with PTB were black (90.9%; 339 cases), 6.4% (24 cases) were white and 2.7% (10 cases) were coloured.



**FIG 3-1 DISTRIBUTION OF ACTIVE TB BY SITE (1995)**

The distribution of active PTB cases by commodity is shown in Table 3-1. The rates in this and subsequent tables and figures are expressed per 1000. The majority of active PTB cases (86.9%) came from the gold mining industry (77.2% of all autopsy cases came from that commodity).

**TABLE 3-1 NUMBER OF CASES AND PREVALENCE OF ACTIVE PTB BY COMMODITY AND POPULATION GROUP (1995)**

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	304	131	20	26	0	-	324	105
Platinum	15	60	1	23	0	-	16	54
Coal	10	83	1	14	0	-	11	56
Asbestos	3	100	1	38	9	122	13	100
Copper	0	-	0	-	1	1 000	1	16
Other	0	-	1	37	0	-	1	23
Unknown	7	81	0	-	0	-	7	48
<b>Total</b>	<b>339</b>	<b>120</b>	<b>24</b>	<b>23</b>	<b>10</b>	<b>101</b>	<b>373</b>	<b>93</b>

The age distribution of cases with active PTB is shown in Table 3-2. Most of the cases were aged between 20 and 69 years (297 cases; 79.6%).

**TABLE 3-2 NUMBER OF CASES AND PREVALENCE OF ACTIVE PTB BY AGE AND POPULATION GROUP (1995)**

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	21	76	1	26	0	-	22	69
30-39	85	95	1	13	0	-	86	87
40-49	101	130	6	52	2	200	109	121
50-59	62	134	3	16	3	167	68	101
60-69	6	103	5	19	1	32	12	34
70-79	0	-	4	16	3	176	7	25
80+	0	-	2	20	1	67	3	25
Unknown	64	187	2	125	0	-	66	179
<b>Total</b>	<b>339</b>	<b>120</b>	<b>24</b>	<b>23</b>	<b>10</b>	<b>101</b>	<b>373</b>	<b>93</b>

## SECTION 4 – SILICOSIS

*Silicotic nodules were found in the lungs of 727 cases (18.2% of all autopsies), 93.7% of which came from the gold mining industry. Of all cases of silicosis, occasional silicotic nodules were found in 24.8% of cases, a few in 20.1%, a moderate number in 38.2% and a large number in 16.9%.*

*The distribution of cases with silicosis by commodity and population group is presented in Table 4-1.*

**TABLE 4-1 NUMBER OF CASES AND PREVALENCE OF SILICOSIS BY COMMODITY AND POPULATION GROUP (1995)**

Commodity	Black		White		Coloured		Unknown		Total	
	N	Rate	N	Rate	N	Rate	N	Rate	N	Rate
Gold	488	211	192	249	0	-	1	500	681	220
Platinum	10	40	4	91	0	-	0	-	14	47
Coal	5	41	3	41	0	-	0	-	8	41
Asbestos	2	67	1	38	2	27	0	-	5	38
Copper	0	-	5	185	1	67	0	-	6	140
Unknown	5	58	7	143	0	-	1	111	13	88
<b>Total</b>	<b>510</b>	<b>180</b>	<b>212</b>	<b>200</b>	<b>3</b>	<b>30</b>	<b>2</b>	<b>167</b>	<b>727</b>	<b>182</b>

*Although the silicosis rates increased with increasing age in both black and white men, the age distribution of cases with silicosis differed between the two population groups (Table 4-2).*

**TABLE 4-2 NUMBER OF CASES AND PREVALENCE OF SILICOSIS IN THE GOLD MINING INDUSTRY, BY AGE AND POPULATION GROUP (1995)**

Age group (years)	Black		White		Coloured		Unknown		Total	
	N	Rate	N	Rate	N	Rate	N	Rate	N	Rate
20-29	6	26	0	-	0	-	0	-	6	23
30-39	74	100	1	19	0	-	0	-	75	95
40-49	180	283	12	143	0	-	1	1000	193	267
50-59	131	350	21	164	0	-	0	-	152	302
60-69	9	281	59	301	0	-	0	-	68	297
70-79	2	333	70	355	0	-	0	-	72	355
80+	0	-	27	321	0	-	0	-	27	300
Unknown	86	306	2	286	0	-	0	-	88	306
<b>Total</b>	<b>488</b>	<b>211</b>	<b>192</b>	<b>249</b>	<b>0</b>		<b>1</b>	<b>500</b>	<b>681</b>	<b>220</b>



Silicosis was diagnosed in men who were young (<40 years) and in men who were exposed to silica for a few years (< 10 years) (Table 4-3).

**TABLE 4-3 NUMBER OF CASES AND PREVALENCE OF SILICOSIS IN THE GOLD MINING INDUSTRY, BY YEARS OF SERVICE AND POPULATION GROUP (1995)**

Years of service	Black		White		Coloured		Unknown		Total	
	N	Rate	N	Rate	N	Rate	N	Rate	N	Rate
<1	4	108	0	-	0	-	0	-	4	105
1-5	34	130	2	31	0	-	0	-	36	110
6-10	43	106	7	99	0	-	0	-	50	105
11-15	94	213	11	151	0	-	0	-	105	203
16-20	123	359	23	274	0	-	1	1000	147	342
21-25	43	457	37	343	0	-	0	-	80	396
26-30	23	451	42	424	0	-	0	-	65	433
31-35	4	235	32	340	0	-	0	-	36	324
36-40	4	444	19	339	0	-	0	-	23	354
41+	0	-	10	435	0	-	0	-	10	435
Unknown	116	177	9	93	0	-	0	-	125	166
<i>Total</i>	488	211	192	249	0		1	500	681	220

## SECTION 5 – OTHER PNEUMOCONIOSES

---

### MASSIVE FIBROSIS

There were 38 (0.9%) cases of massive fibrosis (27 black and 11 white). Thirty five cases of massive fibrosis were from gold, one from platinum and for two cases the commodity was not known.

### COAL WORKERS' PNEUMOCONIOSIS

There were 13 (0.3%) cases of coal workers' pneumoconiosis of which 2 cases were from gold, 10 from coal and for one case the commodity was not known.

### MIXED DUST PNEUMOCONIOSIS

There were 39 (1.0%) cases of mixed dust pneumoconiosis. These cases came from the gold (n=32), coal (n=2) and platinum (2) mining industries. Two cases were from Iscor and for one case the commodity was not known.

### ASBESTOSIS AND PLEURAL PLAQUES

There were 104 cases of asbestosis of which 48.1% (n=50) had slight, 45.2% (n=47) moderate and 6.7% (n=7) marked fibrosis. Of these, 74 (71.1%) had worked in the asbestos mining industry at some time in their lives. There were 37 cases that had asbestos plaques and 20 (54.1%) of these had asbestosis. Note that the parietal pleura (the site where plaque formation usually occurs) is seldom submitted with the lungs.

The distribution of asbestosis by age and population group is shown in Table 5-1.

**TABLE 5-1 NUMBER OF CASES AND PREVALENCE OF ASBESTOSIS BY AGE AND POPULATION GROUP (1995)**

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
30-39	5	6	0	-	0	-	5	5
40-49	6	8	1	9	1	100	8	9
50-59	15	32	7	36	6	333	28	42
60-69	9	155	10	37	14	452	33	93
70-79	1	143	7	28	12	706	20	72
80+	0	-	0	-	5	333	5	41
Unknown	5	15	0	-	0	-	5	14
<b>Total</b>	<b>41</b>	<b>14</b>	<b>25</b>	<b>24</b>	<b>38</b>	<b>384</b>	<b>104</b>	<b>26</b>

## SECTION 6 – EMPHYSEMA

There were 668 cases of emphysema, the extent of which was mild in 59.6% (n=398), moderate in 33.1% (n=221) and marked in 7.3% (n=49). The distribution of emphysema by age and population group is presented in Table 6-1.

**TABLE 6-1 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY AGE AND POPULATION GROUP (1995)**

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	1	4	0	-	1	500	2	6
30-39	38	42	6	75	3	600	47	48
40-49	71	92	23	200	6	600	100	111
50-59	75	162	64	333	11	611	150	223
60-69	13	224	138	517	10	323	161	452
70-79	3	429	126	500	4	235	133	482
80+	2	333	41	410	0	-	43	355
Unknown	31	91	1	63	0	-	32	87
<b>Total</b>	<b>234</b>	<b>83</b>	<b>399</b>	<b>376</b>	<b>35</b>	<b>354</b>	<b>668</b>	<b>167</b>

The majority of black and white men with emphysema were from the gold mining industry (71.1%, n=475) (Table 6-2).

**TABLE 6-2 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY COMMODITY AND POPULATION GROUP (1995)**

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	169	73	306	397	0	-	475	154
Platinum	22	88	15	341	0	-	37	125
Coal	16	132	20	274	0	-	36	185
Asbestos	10	333	12	462	27	365	49	377
Isacor	3	250	19	380	0	-	22	349
Diamond	2	182	6	300	1	1000	9	281
Copper	0	-	4	148	6	400	10	233
Other	2	400	0	-	0	-	2	286
Unknown	10	116	17	347	1	333	28	190
<b>Total</b>	<b>234</b>	<b>83</b>	<b>399</b>	<b>376</b>	<b>35</b>	<b>354</b>	<b>668</b>	<b>167</b>

**TABLE 6-3 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY YEARS OF SERVICE AND POPULATION GROUP (1995)**

Years of service	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
<1	2	50	0	-	0	-	2	48
1-5	17	59	27	267	3	200	47	116
6-10	30	66	33	330	7	304	70	122
11-15	32	69	37	301	8	471	77	127
16-20	33	90	54	443	5	357	92	183
21-25	16	163	60	455	2	333	78	331
26-30	7	135	56	479	2	400	65	374
31-35	3	167	46	447	0	-	49	405
36-40	3	250	28	444	0	-	31	397
41+	0	-	13	500	0	-	13	481
Unknown	91	88	45	260	8	533	144	117
<i>Total</i>	234	83	399	376	35	354	668	167

## SECTION 7 – MESOTHELIOMA

*There were 14 cases of mesothelioma in 1995.*

**TABLE 7-1 NUMBER AND PROPORTION OF MESOTHELIOMA CASES BY AGE AND POPULATION GROUP (1995)**

Age group (years)	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
30-39	1	14.3	0	-	0	-	1	7.1
40-49	4	57.1	0	-	0	-	4	28.6
50-59	2	28.6	0	-	2	66.7	4	28.6
60-69	0	-	2	50.0	1	33.3	3	21.4
70-79	0	-	2	50.0	0	-	2	14.3
<i>Total</i>	7		4		3		14	

*The distribution of mesothelioma by commodity and population group is presented in Table 7.2. Six (50.0%) had worked in the asbestos mining industry.*

**TABLE 7-2 NUMBER AND PROPORTION OF MESOTHELIOMA CASES BY COMMODITY AND POPULATION GROUP (1995)**

Commodity	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
<i>Gold</i>	5	71.4	0	-	0	-	5	35.7
<i>Asbestos</i>	1	14.3	2	50.0	3	100.0	6	42.9
<i>Unknown</i>	1	14.3	2	50.0	0	-	3	21.4
<i>Total</i>	7		4		3		14	

## SECTION 8 – PRIMARY LUNG CANCER

One hundred and three cases of primary lung cancer were found at autopsy, 33.0% of which were in black, 61.2% in white and 5.8% in coloured men. Most of the cases were squamous lung carcinomas (34.9%; n=36), followed by adeno carcinoma (20.4%; n=21), small cell lung carcinoma (19.4%; n=20), large cell lung carcinoma (17.5%; n=18) and broncho-alveolar carcinoma (7.8%; n=8).

The distribution of primary lung cancer by age and population group is presented in Table 8-1.

**TABLE 8-1 NUMBERS OF CASES AND PREVALENCE OF PRIMARY LUNG CANCER BY AGE AND POPULATION GROUP (1995)**

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
30-39	4	4	1	13	0	-	5	5
40-49	10	13	2	17	0	-	12	13
50-59	13	28	15	78	0	-	28	42
60-69	2	34	22	82	5	161	29	81
70-79	0	-	20	79	0	-	20	72
80+	0	-	3	30	1	67	4	33
Unknown	5	15	0	-	0	-	5	14
<b>Total</b>	<b>34</b>	<b>12</b>	<b>63</b>	<b>59</b>	<b>6</b>	<b>61</b>	<b>103</b>	<b>26</b>

The distribution of primary lung cancer by commodity and population group is presented in Table 8-2. The majority of cases had worked in the gold mining industry.

**TABLE 8-2 NUMBER OF CASES AND PREVALENCE OF PRIMARY LUNG CANCER BY COMMODITY AND POPULATION GROUP (1995)**


Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	23	10	45	58	0	-	68	22
Platinum	4	16	2	45	0	-	6	20
Coal	5	41	0	-	0	-	5	26
Asbestos	2	67	4	154	5	68	11	85
Iscor	0	-	5	100	0	-	5	79
Diamond	0	-	1	50	1	1000	2	63
Copper	0	-	1	37	0	-	1	23
Other	0	-	1	500	0	-	1	143
Unknown	0	-	4	82	0	-	4	27
<b>Total</b>	<b>34</b>	<b>12</b>	<b>63</b>	<b>59</b>	<b>6</b>	<b>61</b>	<b>103</b>	<b>26</b>

## SECTION 9 – CLINICAL CAUSES OF DEATH

Table 9-1 and Figure 9-1 show the clinical causes of death as stated in the accompanying documents submitted with the cardio-respiratory organs, by population group. Diseases of the respiratory system were the most frequent (11.6%) overall. Black men had the highest proportion of unnatural causes of death (37.0%). In 35.4% of all cases, the cause of death was not stated.

**TABLE 9-1 CLINICAL CAUSES OF DEATH BY POPULATION GROUP (1995)**

System	Black		White		Coloured		Unknown		Total	
	N	%	N	%	N	%	N	%	N	%
Respiratory	349	12.3	96	9.0	18	18.2	0	-	463	11.6
Cardio-vascular	106	3.7	160	15.1	10	10.1	1	8.3	277	6.9
Central Nervous System	99	3.5	34	3.2	7	7.1	0	-	140	3.5
Gastro-intestinal	97	3.4	39	3.7	6	6.1	0	-	142	3.5
Genito-urinary	39	1.4	24	2.3	0	-	0	-	63	1.6
Haematological	13	0.5	7	0.7	0	-	0	-	20	0.5
Unnatural	1 047	37.0	125	11.8	6	6.1	2	16.7	1 180	29.5
Miscellaneous	207	7.3	84	7.9	8	8.1	1	8.3	300	7.5
Not stated	873	30.8	493	46.4	44	44.4	8	66.7	1 418	35.4
<b>Total</b>	<b>2 830</b>		<b>1 062</b>		<b>99</b>		<b>12</b>		<b>4 003</b>	



**FIGURE 9-1 CLINICAL CAUSE OF DEATH AS STATED BY THE CLINICIANS WHO SUBMIT THE ORGANS OF THE DECEASED TO THE NIOH (1995)**