



**NATIONAL HEALTH
LABORATORY SERVICE**

NATIONAL INSTITUTE FOR OCCUPATIONAL HEALTH

Pathology Division Surveillance Report

Demographic Data and Disease Rates for January to December 1984

Ntombizodwa Ndlovu

Tony Davies

Gill Nelson

Jill Murray

*PO Box 4788
Johannesburg
2000*

e-mail: jill.murray@nioh.nhls.ac.za

NIOH Report 17/2011

http://www.nioh.ac.za/publications/publications_pathaut_reports.htm

ACKNOWLEDGEMENTS

We thank all the staff members of the NIOH Pathology Department, for their invaluable contribution to the autopsy service.

We also thank Prof Tony Davies for editing the manuscript.

National Institute for Occupational Health, PO Box 4788, Johannesburg, 2000, South Africa

ISSN 1812-7681

EXECUTIVE SUMMARY

During 1984, 3 092 cases came to autopsy at the NIOH. Of these, 63.6% were black men, 35.5% were white and 0.9% were coloured.

Overall disease rates (per 1000 autopsies) for 1984 are shown in Figure 1.




FIG 1 OVERALL DISEASE RATES FOR 1984

TABLE OF CONTENTS

EXECUTIVE SUMMARY	<i>i</i>
GLOSSARY	<i>iv</i>
SECTION 1: BACKGROUND	1
SECTION 2: DEMOGRAPHIC DATA	2
SECTION 3: ACTIVE TUBERCULOSIS	7
SECTION 4: SILICOSIS	9
SECTION 5: OTHER PNEUMOCONIOSES	
<i>Massive Fibrosis</i>	11
<i>Coal Workers' Pneumoconiosis</i>	11
<i>Mixed Dust Pneumoconiosis</i>	11
<i>Asbestosis and pleural plaques</i>	11
SECTION 6: EMPHYSEMA	12
SECTION 7: MESOTHELIOMA	14
SECTION 8: PRIMARY LUNG CANCER	15
SECTION 9: CLINICAL CAUSES OF DEATH	16

LIST OF TABLES

Table 2.1	<i>Distribution of autopsies by year and population group (1984)</i>	2
Table 2.2	<i>Number and proportion of autopsies by type and population group (1984)</i>	3
Table 2.3	<i>Number and proportion of autopsies by age and population group (1984)</i>	3
Table 2.4	<i>Number and proportion of autopsies by commodity and population group (1984)</i>	4
Table 2.5	<i>Number and proportion of autopsies by years of service and population group (1984)</i>	5
Table 2.6	<i>Mean age by commodity and population group (1984)</i>	6
Table 2.7	<i>Mean duration of service by commodity and population group (1984)</i>	6
Table 3.1	<i>Number of cases and prevalence of active PTB by commodity and population group (1984)</i>	8
Table 3.2	<i>Number of cases and prevalence of active PTB by age and population group (1984)</i>	8
Table 4.1	<i>Number of cases and prevalence of silicosis by commodity and population group (1984)</i>	9
Table 4.2	<i>Number of cases and prevalence of silicosis in the gold mining industry, by age and population group (1984)</i>	9
Table 4.3	<i>Number of cases and prevalence of silicosis in the gold mining industry, by years of service and population group (1984)</i>	10
Table 5.1	<i>Number of cases and prevalence of asbestosis by age and population group (1984)</i>	11

Table 6.1	<i>Number of cases and prevalence of emphysema by age and population group (1984)</i>	12
Table 6.2	<i>Number of cases and prevalence of emphysema by commodity and population group (1984)</i>	12
Table 6.3	<i>Number of cases and prevalence of emphysema by years of service and population group (1984)</i>	13
Table 7.1	<i>Number and proportion of mesothelioma cases by age and population group (1984)</i>	14
Table 7.2	<i>Number and proportion of mesothelioma cases by commodity and population group (1984)</i>	14
Table 8.1	<i>Number of cases and prevalence of primary lung cancer by age and population group (1984)</i>	15
Table 8.2	<i>Number of cases and prevalence of primary lung cancer by commodity and population group (1984)</i>	15
Table 9.1	<i>Clinical causes of death by population group (1984)</i>	16

LIST OF FIGURES

Figure 1	<i>Overall disease rates for 1984</i>	<i>i</i>
Figure 2.1	<i>Distribution of autopsies by age and population group (1984)</i>	3
Figure 2.2	<i>Distribution of autopsies by commodity and population group (1984)</i>	4
Figure 2.3	<i>Distribution of autopsies by years of service and population group (1984)</i>	5
Figure 3.1	<i>Distribution of active TB by site (1984)</i>	7
Figure 9.1	<i>Clinical cause of death as given by the clinicians who submit the organs to the NIOH (1984)</i>	16

GLOSSARY

Asbestosis	<i>Lung fibrosis caused by asbestos fibres</i>
Coal workers' pneumoconiosis	<i>Lung fibrosis caused by exposure to coal dust</i>
Emphysema	<i>Lung disease caused by the destruction of the alveolar walls</i>
Massive fibrosis	<i>Lung fibrosis caused by exposure to dust and measuring more than 1 cm in diameter</i>
Mesothelioma	<i>A malignant tumour of the pleural cavity of the lungs</i>
Miner	<i>A person who has worked in a controlled mine or works</i>
Mixed exposures	<i>The multiple dust types to which a miner may be exposed, having worked in several mining commodities in his lifetime</i>
Prevalence	<i>The number of cases in a defined population at a given time</i>
Silicosis	<i>Lung fibrosis caused by inhalation of silica dust; detected by the presence of silicotic nodules in the lung tissue</i>
Surveillance	<i>The ongoing and systematic collection, analysis and interpretation of data related to adverse health outcomes</i>

SECTION 1 – BACKGROUND

The Occupational Diseases in Mines and Works Act, 1973 (Act 78 of 1973) requires that the cardio-respiratory organs of a deceased person who has worked at a controlled mine or a controlled works be examined for the presence of occupational disease, regardless of the clinical cause of death and provided that the next of kin agrees. These examinations are performed by pathologists at the National Institute for Occupational Health (NIOH). A detailed report on each case examined is sent to the Medical Bureau for Occupational Diseases (MBOD). Cases certified as having a compensable disease are then referred to the Compensation Commissioner's office, where the payment for compensation is managed.

Since 1975, the pathological findings from the autopsy examinations have been recorded on the computerised PATHAUT database. PATHAUT comprises data from autopsy examinations and clinical files which include occupational histories. The database is unique and provides an important resource for both surveillance and research. These data are the only comprehensive surveillance data on occupational lung disease in the South African mining industry.

The data presented in this report summarise the PATHAUT system surveillance results, i.e. the results of the systematic collection, collation, and analysis of the pathology findings in the cardio-respiratory organs of mine workers. Data from PATHAUT are exported into, and analysed, using SAS v9.1.

This report describes autopsy cases examined during the year 1984. This and other annual reports can be accessed at www.nioh.ac.za.

Many of the cases had "mixed" exposures in that they had been employed in more than one commodity. For simplicity, cases are categorised according to the commodity in which most years of service were recorded, i.e. the commodity in which the miners had worked for the longest period.

All disease rates reported in this document are expressed per 1000. In all calculations, the denominators used are the total numbers of autopsies in specific commodities, age groups or population groups. Some of these rates must be viewed with caution, as the denominators are very small. This applies, for example, to those commodities where few workers are employed (such as manganese mining), and to the older age groups in some instances.

SECTION 2 – DEMOGRAPHIC DATA

The number of autopsies performed for 1984 is presented in Table 2-1.

TABLE 2-1 DISTRIBUTION OF AUTOPSIES BY YEAR AND POPULATION GROUP (1984)

Year of autopsy	Black		White		Coloured		Unknown		Total N
	N	%	N	%	N	%	N	%	
1975	2 190	71	854	28	32	1			3 076
1976	2 335	68	1 072	31	27	1			3 434
1977	2 351	69	1 039	30	33	1			3 423
1978	2 245	67	1 090	32	32	1			3 367
1979	2 118	66	1 026	33	45	1			3 189
1980	2 338	64	1 274	35	46	1			3 658
1981	2 209	66	1 117	33	33	1			3 359
1982	2 312	63	1 302	36	44	1	1		3 659
1983	2 096	65	1 109	34	41	1			3 246
1984	1 966	64	1 098	36	28	1			3 092
Total	22 160	66	10 981	33	361	1	1		33 503

It is important to note that a referral bias exists: there is a low autopsy rate for black men who have left employment at the mines, whereas the majority of white retired miners come to autopsy.

The pathologists at the NIOH perform two types of autopsy examinations. For men dying distant from Johannesburg, the cardio-respiratory organs are removed locally, preserved in formalin and sent to the NIOH. Full autopsies are undertaken on men who die close to Johannesburg.

Table 2-2 shows the distribution of autopsies by population group for 1984. Autopsies of only the cardio-respiratory organs comprised 83.2% of all examinations.

TABLE 2-2 NUMBER AND PROPORTION OF AUTOPSIES BY TYPE AND POPULATION GROUP (1984)

Autopsy type	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
Cardio-respiratory organs only	1 837	93.4	707	64.4	28	100	2 572	83.2
Full autopsy	129	6.6	391	35.6	0	-	520	16.8
Total	1 966		1 098		28		3 092	

The age distributions of autopsies for 1984 are shown in Table 2-3 and Figure 2-1. The mean age at autopsy of black men was 35.2 years. The mean age of white men at autopsy was 58.1 years and for coloured men 54.4 years

TABLE 2-3 NUMBER AND PROPORTION OF AUTOPSIES BY AGE AND POPULATION GROUP (1984)

Age group (years)	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
<20	23	1.2	3	0.3	0	-	26	0.8
20-29	634	32.2	56	5.1	1	3.6	691	22.3
30-39	566	28.8	74	6.7	2	7.1	642	20.8
40-49	386	19.6	118	10.7	7	25.0	511	16.5
50-59	215	10.9	218	19.9	6	21.4	439	14.2
60-69	70	3.6	329	30.0	9	32.1	408	13.2
70-79	9	0.5	241	21.9	2	7.1	252	8.2
80+	0	-	40	3.6	1	3.6	41	1.3
Unknown	63	3.2	19	1.7	0	-	82	2.7
Total	1 966		1 098		28		3 092	




FIG 2-1 DISTRIBUTION OF AUTOPSIES BY AGE AND POPULATION GROUP (1984)

Cases were placed in categories according to the commodity in which they had worked for the longest duration (most exposure). Many men worked in a number of different mining commodities

during their lifetimes and had “mixed” exposures. This was not taken into account in the analysis of exposure type (commodity).

Table 2-4 and Figure 2-2 show the distribution of autopsies by commodity and population group for 1984.

TABLE 2-4 NUMBER AND PROPORTION OF AUTOPSIES BY COMMODITY AND POPULATION GROUP (1984)

Commodity	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
Gold	1 558	79.2	807	73.6	4	14.3	2 369	76.6
Platinum	120	6.1	26	2.4	0	-	146	4.7
Coal	198	10.1	68	6.2	0	-	266	8.6
Asbestos	15	0.8	20	1.8	20	71.4	55	1.8
Iscor	10	0.5	87	7.9	1	3.6	98	3.2
Diamond	7	0.4	9	0.8	0	-	16	0.5
Copper	10	0.5	21	1.9	3	10.7	34	1.1
Other	5	0.3	2	0.2	0	-	7	0.2
Unknown	43	2.2	58	5.3	0	-	101	3.3
Total	1 966		1 098		28		3 092	

NOTE: This table shows only those commodities where a total of 10 or more cases were received





FIG 2-2 DISTRIBUTION OF AUTOPSIES BY COMMODITY AND POPULATION GROUP (1984)

Detailed information about the years in mining service by population group is presented in Table 2-5 and Figure 2-3.

TABLE 2-5 NUMBER AND PROPORTION OF AUTOPSIES BY YEARS OF SERVICE AND POPULATION GROUP (1984)

Years of service	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
<1	160	8.1	10	0.9	1	3.6	171	5.5
1-5	655	33.3	83	7.6	5	17.9	743	24.0
6-10	379	19.3	77	7.0	9	32.1	465	15.0
11-15	150	7.6	91	8.3	3	10.7	244	7.9
16-20	76	3.9	98	8.9	3	10.7	177	5.7
21-25	28	1.4	108	9.8	1	3.6	137	4.4
26-30	17	0.9	141	12.8	0	-	158	5.1
31-35	8	0.4	142	12.9	1	3.6	151	4.9
36-40	8	0.4	118	10.7	0	-	126	4.1
41+	1	0.1	70	6.4	2	7.1	73	2.4
Unknown	484	24.6	160	14.6	3	10.7	647	20.9
Total	1 966		1 098		28		3 092	



Years of service:




FIG 2-3 DISTRIBUTION OF AUTOPSIES BY YEARS OF SERVICE AND POPULATION GROUP (1984)

The mean age and duration of service by commodity type and population group for those cases for which information was available are shown in Tables 2-6 and 2-7.

TABLE 2-6 MEAN AGE BY COMMODITY AND POPULATION GROUP (1984)

Commodity	Black			White		
	N	Mean (years)	SD*	N	Mean (years)	SD*
Gold	1501	36	11	806	60	15
Platinum	119	36	11	26	51	16
Coal	196	37	13	68	55	17
Asbestos	15	43	11	20	65	13
Iscor	10	41	15	87	56	13
Diamond	7	40	11	9	54	13
Copper	9	49	9	21	55	16
Other	5	46	11	2	47	19
Unknown	41	43	13	40	60	14
Total	1 903	36	12	1 079	59	15

* Standard deviation

TABLE 2-7 MEAN DURATION OF SERVICE BY COMMODITY AND POPULATION GROUP (1984)

Commodity	Black			White		
	N	Mean (years)	SD*	N	Mean (years)	SD*
Gold	1230	7	6	742	25	12
Platinum	90	8	3	23	13	7
Coal	120	8	9	56	20	13
Asbestos	12	12	10	14	15	9
Iscor	5	15	15	56	22	12
Diamond	5	16	10	9	16	13
Copper	4	9	9	20	19	12
Other	3	14	11	2	18	16
Unknown	13	9	5	16	18	5
Total	1 482	7	7	938	24	13

* Standard deviation

SECTION 3 – ACTIVE TUBERCULOSIS

The distribution of active tuberculosis (TB) by anatomical site is presented in Figure 3-1 (n=172). Active pulmonary TB (PTB) was diagnosed in 4.2% (n=130) of all cases autopsied in 1984. Most of the men with PTB were black (85.4%; 111 cases), 12.3% (16 cases) were white and 2.3% (3 cases) were coloured.




FIG 3-1 DISTRIBUTION OF ACTIVE TB BY SITE (1984)

The distribution of active PTB cases by commodity is shown in Table 3-1. The rates in this and subsequent tables and figures are expressed per 1000. The majority of active PTB cases (53.4%) came from the gold mining industry (71.4% of all autopsy cases came from that commodity).

TABLE 3-1 NUMBER OF CASES AND PREVALENCE OF ACTIVE PTB BY COMMODITY AND POPULATION GROUP (1984)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	90	58	15	41	2	500	107	45
Platinum	8	67	0	-	0	-	8	55
Coal	8	40	0	-	0	-	8	30
Asbestos	1	67	1	43	1	50	3	55
Other	1	200	0	-	0	-	1	143
Unknown	3	70	0	-	0	-	3	30
Total	111	359	16	15	3	107	130	42

The age distribution of cases with active PTB is shown in Table 3-2. Most of the cases were between 20-69 years (119 cases=91.5%).

TABLE 3-2 NUMBER OF CASES AND PREVALENCE OF ACTIVE PTB BY AGE AND POPULATION GROUP (1984)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
<20	1	43	0	-	0	-	1	38
20-29	16	27	0	-	0	-	16	23
30-39	23	41	0	-	1	500	24	37
40-49	36	93	1	8	1	143	38	74
50-59	26	121	0	-	0	-	26	59
60-69	7	100	6	18	1	111	14	34
70-79	0	-	7	29	0	-	7	28
80+	0	-	2	50	0	-	2	49
Unknown	2	32	0	-	0	-	2	24
Total	111	56	16	15	3	107	130	42

SECTION 4 – SILICOSIS

Silicotic nodules were found in the lungs of 391 cases (12.6% of all autopsies), 91.0% of which came from the gold mining industry. Of all cases of silicosis, occasional silicotic nodules were found in 48.8% of cases, a few in 26.5%, a moderate number in 18.4% and a large number in 6.2%.

The distribution of cases with silicosis by commodity and population group is presented in Table 4-1.

TABLE 4-1 NUMBER OF CASES AND PREVALENCE OF SILICOSIS BY COMMODITY AND POPULATION GROUP (1984)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	135	87	221	274	0	-	356	150
Platinum	3	25	1	38	0	-	4	27
Coal	2	10	5	74	0	-	7	26
Asbestos	1	67	2	100	1	50	4	73
Copper	0	-	6	286	0	-	6	176
Isacor	0	-	4	46	0	-	4	41
Unknown	2	47	8	138	0	-	10	99
Total	143	73	247	225	1	36	391	126

Although the silicosis rates increased with increasing age in both black and white men, the age distribution of cases with silicosis differed between the two population groups (Table 4-2).

TABLE 4-2 NUMBER OF CASES AND PREVALENCE OF SILICOSIS IN THE GOLD MINING INDUSTRY, BY AGE AND POPULATION GROUP (1984)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	1	2	0	-	0	-	1	2
30-39	30	65	1	22	0	-	31	61
40-49	61	203	7	91	0	-	68	179
50-59	25	160	31	196	0	-	56	178
60-69	12	235	91	355	0	-	103	334
70-79	0	-	72	369	0	-	72	362
80+	0	-	18	581	0	-	18	581
Unknown	6	105	1	1000	0	-	7	121
Total	135	87	221	274	0	-	356	150

Silicosis was diagnosed in men who were young (<40 years) and in men who were exposed to silica for a few years (< 10 years) (Table 4-3).

TABLE 4-3 NUMBER OF CASES AND PREVALENCE OF SILICOSIS IN THE GOLD MINING INDUSTRY, BY YEARS OF SERVICE AND POPULATION GROUP (1984)

Years of service	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
<1	5	36	0	-	0	-	5	34
1-5	15	28	2	34	0	-	17	28
6-10	29	91	2	41	0	-	31	84
11-15	29	232	9	164	0	-	38	211
16-20	13	200	12	160	0	-	25	179
21-25	3	143	25	294	0	-	28	264
26-30	4	333	41	357	0	-	45	354
31-35	1	167	53	434	0	-	54	422
36-40	0	-	46	422	0	-	46	411
41+	1	1000	26	400	0	-	27	409
Unknown	35	107	5	77	0	-	40	102
<i>Total</i>	135	87	221	274	0	-	356	150

SECTION 5 – OTHER PNEUMOCONIOSES

MASSIVE FIBROSIS

There were 5 (0.2%) cases of massive fibrosis (3 black, 2 white). Four cases of massive fibrosis were from the gold mining industry. The remaining case was from the coal mining industry.

COAL WORKERS' PNEUMOCONIOSIS

There were 25 (0.8%) cases of coal workers' pneumoconiosis of which 23 cases were known to be from the coal mining industry. The remaining 2 cases were from the gold mining industry.

MIXED DUST PNEUMOCONIOSIS

There were 17 (0.5%) cases of mixed dust pneumoconiosis. These cases came from the gold (n=12), coal (n=2), copper (1) and diamond (1) mining industries. In one case the industry was not stated.

ASBESTOSIS AND PLEURAL PLAQUES

There were 66 cases of asbestosis of which 81.8% (n=54) had slight, 13.6% (n=9) moderate and 4.5% (n=3) marked fibrosis. Of these, 36 (54.5%) had worked in the asbestos mining industry at some time in their lives. There were 13 cases that had asbestos plaques and 7 (53.8%) of these had asbestosis. Note that the parietal pleura (the site where plaque formation usually occurs) is seldom submitted with the lungs.

The distribution of asbestosis by age and population group is shown in Table 5-1.

TABLE 5-1 NUMBER OF CASES AND PREVALENCE OF ASBESTOSIS BY AGE AND POPULATION GROUP (1984)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	1	2	0	-	0	-	1	1
30-39	5	9	0	-	0	-	5	8
40-49	13	34	0	-	3	429	16	31
50-59	7	33	4	18	3	500	14	32
60-69	4	57	8	24	6	667	18	44
70-79	0	-	2	8	2	1000	4	16
80+	0	-	4	100	1	1000	5	122
Unknown	1	16	2	105	0	-	3	37
Total	31	16	20	18	15	536	66	21

SECTION 6 – EMPHYSEMA

There were 915 cases of emphysema, the extent of which was mild in 73.2% (n=670), moderate in 24.7% (n=226) and marked in 2.1% (n=19). The distribution of emphysema by age and population group is presented in Table 6-1.

TABLE 6-1 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY AGE AND POPULATION GROUP (1984)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	9	14	0	-	0	-	9	13
30-39	41	72	5	68	1	500	47	73
40-49	93	241	31	263	0	-	124	243
50-59	75	349	141	647	2	333	218	497
60-69	26	371	259	787	8	889	293	718
70-79	5	556	177	734	1	500	183	726
80+	0	-	30	750	0	-	30	732
Unknown	6	95	5	263	0	-	11	134
Total	255	130	648	590	12	429	915	296

The majority of black and white men with emphysema were from the gold mining industry (73.7%, n=674) (Table 6-2).

TABLE 6-2 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY COMMODITY AND POPULATION GROUP (1984)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	173	111	501	621	1	250	675	285
Platinum	14	117	7	269	0	-	21	144
Coal	45	227	32	471	0	-	77	289
Asbestos	2	133	16	800	9	450	27	491
Diamond	3	429	4	444	0	-	7	438
Copper	5	500	11	524	2	667	18	529
Iscor	1	100	49	563	0	-	50	510
Other	2	400	0	-	0	-	2	286
Unknown	10	233	28	483	0	-	38	376
Total	255	130	648	590	12	429	915	296

TABLE 6-3 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY YEARS OF SERVICE AND POPULATION GROUP (1984)

Years of service	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
<1	7	44	0	-	0	-	7	41
1-5	38	58	27	325	3	600	68	92
6-10	44	116	34	442	5	556	83	178
11-15	34	227	45	495	1	333	80	328
16-20	16	211	62	633	1	333	79	446
21-25	8	286	70	648	0	-	78	569
26-30	8	471	93	660	0	-	101	639
31-35	4	500	111	782	0	-	115	762
36-40	4	500	90	763	0	-	94	746
41+	0	-	49	700	2	1000	51	699
Unknown	92	190	67	419	0	-	159	246
<i>Total</i>	255	130	648	590	12	429	915	296

SECTION 7 – MESOTHELIOMA

There were 16 cases of mesothelioma in 1984.

TABLE 7-1 NUMBER AND PROPORTION OF MESOTHELIOMA CASES BY AGE AND POPULATION GROUP (1984)

Age group (years)	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
20-29	1	20.0	0	-	0	-	1	6
30-39	1	20.0	0	-	0	-	1	6
40-49	1	20.0	0	-	2	67	3	19
50-59	2	40.0	0	-	1	33	3	19
60-69	0	-	4	50	0	-	4	25
70-79	0	-	4	50	0	-	4	25
<i>Total</i>	5		8		3		16	

The distribution of mesothelioma by commodity and population group is presented in Table 7.2. Seven (44.0%) had worked in the asbestos mining industry.

TABLE 7-2 NUMBER AND PROPORTION OF MESOTHELIOMA CASES BY COMMODITY AND POPULATION GROUP (1984)

Commodity	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
<i>Gold</i>	1	20	4	50	0	-	5	31
<i>Asbestos</i>	3	60	1	13	3	100	7	44
<i>Iscor</i>	0	-	1	13	0	-	1	6
<i>Unknown</i>	1	20	2	25	0	-	3	19
<i>Total</i>	5		8		3		16	

SECTION 8 – PRIMARY LUNG CANCER

One hundred and thirteen cases of primary lung cancer were found at autopsy, 19.5% of which were in black, 79.6% in white and 0.9% in coloured men. Most of the cases were small cell lung carcinomas (40.7%; n=46), followed by squamous lung carcinomas (29.2%; n=33), adenocarcinomas (15.0%; n=17), large cell lung carcinomas (10.6%; n=12) and broncho-alveolar carcinomas (4.4%; n=5)

The distribution of primary lung cancer by age and population group is presented in Table 8-1.

TABLE 8-1 NUMBERS OF CASES AND PREVALENCE OF PRIMARY LUNG CANCER BY AGE AND POPULATION GROUP (1984)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	1	2	0	-	0	-	1	1
30-39	3	5	0	-	0	-	3	5
40-49	6	16	5	42	0	-	11	22
50-59	10	47	17	78	0	-	27	62
60-69	2	29	34	103	1	111	37	91
70-79	0	-	31	129	0	-	31	123
80+	0	-	2	50	0	-	2	49
Unknown	0	-	1	53	0	-	1	12
Total	22	11	90	82	1	36	113	37

The distribution of primary lung cancer by commodity and population group is presented in Table 8-2. The majority of cases came from the gold mining industry.

TABLE 8-2 NUMBER OF CASES AND PREVALENCE OF PRIMARY LUNG CANCER BY COMMODITY AND POPULATION GROUP (1984)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	14	9	69	86	1	250	84	35
Platinum	3	25	0	-	0	-	3	21
Coal	3	15	3	44	0	-	6	23
Asbestos	1	67	2	100	0	-	3	55
Copper	0	-	1	48	0	-	1	29
Iscor	0	-	5	57	0	-	5	51
Unknown	1	23	10	172	0	-	11	109
Total	22	11	90	82	1	36	113	37

SECTION 9 – CLINICAL CAUSES OF DEATH

Table 9-1 and Figure 9-1 show the clinical causes of death as stated in the accompanying documents submitted with the cardio-respiratory organs, by population group. Diseases of the cardio-vascular system were the most frequent (11.1%) overall. Black men had the highest proportion of unnatural causes of death (64.2%). In 20.7% of all cases, the cause of death was not stated.

TABLE 9-1 CLINICAL CAUSES OF DEATH BY POPULATION GROUP (1984)

System	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
Respiratory	175	8.9	197	17.9	6	21.4	378	12.2
Cardio-vascular	103	5.2	442	40.3	7	25.0	552	17.9
Central Nervous System	86	4.4	51	4.6	3	10.7	140	4.5
Gastro-intestinal	107	5.4	74	6.7	2	7.1	183	5.9
Genito-urinary	24	1.2	35	3.2	1	3.6	60	1.9
Haematological	24	1.2	9	0.8	1	3.6	34	1.1
Unnatural	1 263	64.2	163	14.8	2	7.1	1 428	46.2
Miscellaneous	127	6.5	63	5.7	4	14.3	194	6.3
Not stated	57	2.9	64	5.8	2	7.1	123	4.0
Total	1 966		1 098		28		3 092	




FIGURE 9-1 CLINICAL CAUSE OF DEATH AS STATED BY THE CLINICIANS WHO SUBMIT THE ORGANS OF THE DECEASED TO THE NIOH (1984)