



**NATIONAL HEALTH
LABORATORY SERVICE**

NATIONAL INSTITUTE FOR OCCUPATIONAL HEALTH

Pathology Division Surveillance Report

Demographic Data and Disease Rates for January to December 1986

Ntombizodwa Ndlovu

Tony Davies

Gill Nelson

Jill Murray

*PO Box 4788
Johannesburg
2000*

e-mail: jill.murray@nioh.nhls.ac.za

NIOH Report 19/2011

http://www.nioh.ac.za/publications/publications_pathaut_reports.htm

ACKNOWLEDGEMENTS

We thank all the staff members of the NIOH Pathology Department, for their invaluable contribution to the autopsy service.

We also thank Prof Tony Davies for editing the manuscript.

National Institute for Occupational Health, PO Box 4788, Johannesburg, 2000, South Africa

ISSN 1812-7681

EXECUTIVE SUMMARY

During 1986, 3 626 cases came to autopsy at the NIOH. Of these, 67.7% were black men, 31.0% were white and 1.2% were coloured.

Overall disease rates (per 1000 autopsies) for 1986 are shown in Figure 1.




FIG 1 OVERALL DISEASE RATES FOR 1986

TABLE OF CONTENTS

EXECUTIVE SUMMARY	<i>i</i>
GLOSSARY	<i>iv</i>
SECTION 1: BACKGROUND	1
SECTION 2: DEMOGRAPHIC DATA	2
SECTION 3: ACTIVE TUBERCULOSIS	7
SECTION 4: SILICOSIS	9
SECTION 5: OTHER PNEUMOCONIOSES	
<i>Massive Fibrosis</i>	11
<i>Coal Workers' Pneumoconiosis</i>	11
<i>Mixed Dust Pneumoconiosis</i>	11
<i>Asbestosis and pleural plaques</i>	11
SECTION 6: EMPHYSEMA	12
SECTION 7: MESOTHELIOMA	14
SECTION 8: PRIMARY LUNG CANCER	15
SECTION 9: CLINICAL CAUSES OF DEATH	16

LIST OF TABLES

Table 2.1	<i>Distribution of autopsies by year and population group (1986)</i>	2
Table 2.2	<i>Number and proportion of autopsies by type and population group (1986)</i>	2
Table 2.3	<i>Number and proportion of autopsies by age and population group (1986)</i>	3
Table 2.4	<i>Number and proportion of autopsies by commodity and population group (1986)</i>	4
Table 2.5	<i>Number and proportion of autopsies by years of service and population group (1986)</i>	5
Table 2.6	<i>Mean age by commodity and population group (1986)</i>	6
Table 2.7	<i>Mean duration of service by commodity and population group (1986)</i>	6
Table 3.1	<i>Number of cases and prevalence of active PTB by commodity and population group (1986)</i>	8
Table 3.2	<i>Number of cases and prevalence of active PTB by age and population group (1986)</i>	8
Table 4.1	<i>Number of cases and prevalence of silicosis by commodity and population group (1986)</i>	9
Table 4.2	<i>Number of cases and prevalence of silicosis in the gold mining industry, by age and population group (1986)</i>	9
Table 4.3	<i>Number of cases and prevalence of silicosis in the gold mining industry, by years of service and population group (1986)</i>	10
Table 5.1	<i>Number of cases and prevalence of asbestosis by age and population group (1986)</i>	11

Table 6.1	<i>Number of cases and prevalence of emphysema by age and population group (1986)</i>	12
Table 6.2	<i>Number of cases and prevalence of emphysema by commodity and population group (1986)</i>	12
Table 6.3	<i>Number of cases and prevalence of emphysema by years of service and population group (1986)</i>	13
Table 7.1	<i>Number and proportion of mesothelioma cases by age and population group (1986)</i>	14
Table 7.2	<i>Number and proportion of mesothelioma cases by commodity and population group (1986)</i>	14
Table 8.1	<i>Number of cases and prevalence of primary lung cancer by age and population group (1986)</i>	15
Table 8.2	<i>Number of cases and prevalence of primary lung cancer by commodity and population group (1986)</i>	15
Table 9.1	<i>Clinical causes of death by population group (1986)</i>	16

LIST OF FIGURES

Figure 1	<i>Overall disease rates for 1986</i>	<i>i</i>
Figure 2.1	<i>Distribution of autopsies by age and population group (1986)</i>	3
Figure 2.2	<i>Distribution of autopsies by commodity and population group (1986)</i>	4
Figure 2.3	<i>Distribution of autopsies by years of service and population group (1986)</i>	5
Figure 3.1	<i>Distribution of active TB by site (1986)</i>	7
Figure 9.1	<i>Clinical cause of death as given by the clinicians who submit the organs to the NIOH (1986)</i>	16

GLOSSARY

Asbestosis	<i>Lung fibrosis caused by asbestos fibres</i>
Coal workers' pneumoconiosis	<i>Lung fibrosis caused by exposure to coal dust</i>
Emphysema	<i>Lung disease caused by the destruction of the alveolar walls</i>
Massive fibrosis	<i>Lung fibrosis caused by exposure to dust and measuring more than 1 cm in diameter</i>
Mesothelioma	<i>A malignant tumour of the pleural cavity of the lungs</i>
Miner	<i>A person who has worked in a controlled mine or works</i>
Mixed exposures	<i>The multiple dust types to which a miner may be exposed, having worked in several mining commodities in his lifetime</i>
Prevalence	<i>The number of cases in a defined population at a given time</i>
Silicosis	<i>Lung fibrosis caused by inhalation of silica dust; detected by the presence of silicotic nodules in the lung tissue</i>
Surveillance	<i>The ongoing and systematic collection, analysis and interpretation of data related to adverse health outcomes</i>

SECTION 1 – BACKGROUND

The Occupational Diseases in Mines and Works Act, 1973 (Act 78 of 1973) requires that the cardio-respiratory organs of a deceased person who has worked at a controlled mine or a controlled works be examined for the presence of occupational disease, regardless of the clinical cause of death and provided that the next of kin agrees. These examinations are performed by pathologists at the National Institute for Occupational Health (NIOH). A detailed report on each case examined is sent to the Medical Bureau for Occupational Diseases (MBOD). Cases certified as having a compensable disease are then referred to the Compensation Commissioner's office, where the payment for compensation is managed.

Since 1975, the pathological findings from the autopsy examinations have been recorded on the computerised PATHAUT database. PATHAUT comprises data from autopsy examinations and clinical files which include occupational histories. The database is unique and provides an important resource for both surveillance and research. These data are the only comprehensive surveillance data on occupational lung disease in the South African mining industry.

The data presented in this report summarise the PATHAUT system surveillance results, i.e. the results of the systematic collection, collation, and analysis of the pathology findings in the cardio-respiratory organs of mine workers. Data from PATHAUT are exported into, and analysed, using SAS v9.1.

This report describes autopsy cases examined during the year 1986. This and other annual reports can be accessed at www.nioh.ac.za.

Many of the cases had "mixed" exposures in that they had been employed in more than one commodity. For simplicity, cases are categorised according to the commodity in which most years of service were recorded, i.e. the commodity in which the miners had worked for the longest period.

All disease rates reported in this document are expressed per 1000. In all calculations, the denominators used are the total numbers of autopsies in specific commodities, age groups or population groups. Some of these rates must be viewed with caution, as the denominators are very small. This applies, for example, to those commodities where few workers are employed (such as manganese mining), and to the older age groups in some instances.

SECTION 2 – DEMOGRAPHIC DATA

The number of autopsies performed for 1986 is presented in Table 2-1.

TABLE 2-1 DISTRIBUTION OF AUTOPSIES BY YEAR AND POPULATION GROUP (1986)

Year of autopsy	Black		White		Coloured		Unknown		Total N
	N	%	N	%	N	%	N	%	
1975	2 190	71	854	28	32	1			3 076
1976	2 335	68	1 072	31	27	1			3 434
1977	2 351	69	1 039	30	33	1			3 423
1978	2 245	67	1 090	32	32	1			3 367
1979	2 118	66	1 026	33	45	1			3 189
1980	2 338	64	1 274	35	46	1			3 658
1981	2 209	66	1 117	33	33	1			3 359
1982	2 312	63	1 302	36	44	1	1		3 659
1983	2 096	65	1 109	34	41	1			3 246
1984	1 966	64	1 098	36	28	1			3 092
1985	2 275	64	1 200	34	66	2			3 541
1986	2 456	68	1 125	31	45	1			3 626
Total	26 891	66	13 306	33	472	1	1		40 670

It is important to note that a referral bias exists: there is a low autopsy rate for black men who have left employment at the mines, whereas the majority of white retired miners come to autopsy.

The pathologists at the NIOH perform two types of autopsy examinations. For men dying distant from Johannesburg, the cardio-respiratory organs are removed locally, preserved in formalin and sent to the NIOH. Full autopsies are undertaken on men who die close to Johannesburg.

Table 2-2 shows the distribution of autopsies by population group for 1986. Autopsies of only the cardio-respiratory organs comprised 86.9% of all examinations.

TABLE 2-2 NUMBER AND PROPORTION OF AUTOPSIES BY TYPE AND POPULATION GROUP (1986)

Autopsy type	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
Cardio-respiratory organs only	2 288	93.2	744	66.1	45	100.0	3077	86.9
Full autopsy	140	5.7	370	32.9	0	-	510	14.4
Not stated	28	1.1	11	1.0	0	-	39	1.1
Total	2 456		1 125		45		3 626	

The age distributions of autopsies for 1986 are shown in Table 2-3 and Figure 2-1. The mean age at autopsy of black men was 36.0 years. The mean age of white men at autopsy was 58.9 years and for coloured men 61.8 years

TABLE 2-3 NUMBER AND PROPORTION OF AUTOPSIES BY AGE AND POPULATION GROUP (1986)

Age group (years)	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
<20	8	0.3	5	0.4	0	-	13	0.4
20-29	774	31.5	52	4.6	2	4.4	828	22.8
30-39	694	28.3	56	5.0	3	6.7	753	20.8
40-49	514	20.9	121	10.8	4	8.9	639	17.6
50-59	308	12.5	250	22.2	7	15.6	565	15.6
60-69	73	3.0	328	29.2	16	35.6	417	11.5
70-79	13	0.5	254	22.6	7	15.6	274	7.6
80+	2	0.1	46	4.1	6	13.3	54	1.5
Unknown	70	2.9	13	1.2	0	-	83	2.3
Total	2 456		1 125		45		3 626	




FIG 2-1 DISTRIBUTION OF AUTOPSIES BY AGE AND POPULATION GROUP (1986)

Cases were placed in categories according to the commodity in which they had worked for the longest duration (most exposure). Many men worked in a number of different mining commodities

during their lifetimes and had “mixed” exposures. This was not taken into account in the analysis of exposure type (commodity).

Table 2-4 and Figure 2-2 show the distribution of autopsies by commodity and population group for 1986.

TABLE 2-4 NUMBER AND PROPORTION OF AUTOPSIES BY COMMODITY AND POPULATION GROUP (1986)

Commodity	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
Gold	1 948	79.3	850	75.6	0	-	2 798	77.2
Platinum	182	7.4	29	2.6	0	-	211	5.8
Coal	219	8.9	78	6.9	2	4.4	299	8.2
Asbestos	23	0.9	23	2.0	40	88.9	86	2.4
Iscor	7	0.3	70	6.2	0	-	77	2.1
Diamond	17	0.7	10	0.9	0	-	27	0.7
Copper	6	0.2	17	1.5	2	4.4	25	0.7
Other	8	0.3	7	0.6	0	-	15	0.4
Unknown	46	1.9	41	3.6	1	2.2	88	2.4
Total	2 456		1 125		45		3 626	

NOTE: This table shows only those commodities where a total of 10 or more cases were received




FIG 2-2 DISTRIBUTION OF AUTOPSIES BY COMMODITY AND POPULATION GROUP (1986)

Detailed information about the years in mining service by population group is presented in Table 2-5 and Figure 2-3.

TABLE 2-5 NUMBER AND PROPORTION OF AUTOPSIES BY YEARS OF SERVICE AND POPULATION GROUP (1986)

Years of service	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
<1	185	7.5	5	0.4	0	-	190	5.2
1-5	756	30.8	81	7.2	7	15.6	844	23.3
6-10	539	21.9	81	7.2	12	26.7	632	17.4
11-15	196	8.0	92	8.2	5	11.1	293	8.1
16-20	97	3.9	109	9.7	6	13.3	212	5.8
21-25	36	1.5	126	11.2	3	6.7	165	4.6
26-30	25	1.0	146	13.0	2	4.4	173	4.8
31-35	6	0.2	161	14.3	2	4.4	169	4.7
36-40	2	0.1	145	12.9	1	2.2	148	4.1
41+	2	0.1	64	5.7	1	2.2	67	1.8
Unknown	612	24.9	115	10.2	6	13.3	733	20.2
Total	2 456		1 125		45		3 626	




FIG 2-3 DISTRIBUTION OF AUTOPSIES BY YEARS OF SERVICE AND POPULATION GROUP (1986)

The mean age and duration of service by commodity type and population group for those cases for which information was available are shown in Tables 2-6 and 2-7.

TABLE 2-6 MEAN AGE BY COMMODITY AND POPULATION GROUP (1986)

Commodity	Black			White		
	N	Mean (years)	SD*	N	Mean (years)	SD*
Gold	1 899	36	11	844	60	15
Platinum	178	36	11	29	52	16
Coal	211	40	13	78	60	14
Asbestos	22	49	18	23	55	11
Iscor	7	51	6	70	57	12
Diamond	15	47	10	10	55	13
Copper	5	36	9	16	58	13
Other	8	50	10	7	61	9
Unknown	41	46	12	35	57	11
Total	2 386	37	11	1 112	60	14

* Standard deviation

TABLE 2-7 MEAN DURATION OF SERVICE BY COMMODITY AND POPULATION GROUP (1986)

Commodity	Black			White		
	N	Mean (years)	SD*	N	Mean (years)	SD*
Gold	1 499	7	6	793	26	12
Platinum	142	5	4	25	14	9
Coal	151	10	8	69	23	13
Asbestos	15	8	5	19	14	11
Iscor	3	9	13	59	20	11
Diamond	6	10	8	9	20	10
Copper	2	3	0	14	20	9
Other	7	16	7	7	18	13
Unknown	19	8	5	15	22	11
Total	1 844	7	6	1 010	25	12

* Standard deviation

SECTION 3 – ACTIVE TUBERCULOSIS

The distribution of active tuberculosis (TB) by anatomical site is presented in Figure 3-1 (n=189). Active pulmonary TB (PTB) was diagnosed in 3.8% (n=139) of all cases autopsied in 1986. Most of the men with PTB were black (82.7%; 115 cases), 12.9% (18 cases) were white and 4.3% (6 cases) were coloured.




FIG 3-1 DISTRIBUTION OF ACTIVE TB BY SITE (1986)

The distribution of active PTB cases by commodity is shown in Table 3-1. The rates in this and subsequent tables and figures are expressed per 1000. The majority of active PTB cases (77.0%) came from the gold mining industry (77.2% of all autopsy cases came from that commodity).

TABLE 3-1 NUMBER OF CASES AND PREVALENCE OF ACTIVE PTB BY COMMODITY AND POPULATION GROUP (1986)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	94	48	13	15	0	-	107	38
Platinum	8	44	0	-	0	-	8	38
Coal	9	41	2	26	0	-	11	37
Asbestos	2	87	1	43	6	150	9	105
Iscor	1	143	0	-	0	-	1	13
Copper	0	-	1	59	0	-	1	40
Unknown	1	22	1	24	0	-	2	23
Total	115	47	18	16	6	133	139	38

The age distribution of cases with active PTB is shown in Table 3-2. Most of the cases were aged between 20 and 69 years (134 cases=96.4%).

TABLE 3-2 NUMBER OF CASES AND PREVALENCE OF ACTIVE PTB BY AGE AND POPULATION GROUP (1986)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	14	18	0	-	0	-	14	17
30-39	25	36	0	-	1	333	26	35
40-49	32	62	1	8	0	-	33	52
50-59	32	104	4	16	0	-	36	64
60-69	6	82	4	12	4	250	14	34
70-79	3	231	8	31	0	-	11	40
80+	0	-	0	-	1	167	1	19
Unknown	3	43	1	77	0	-	4	48
Total	115	47	18	16	6	133	139	38

SECTION 4 – SILICOSIS

Silicotic nodules were found in the lungs of 417 cases (11.5% of all autopsies), 93.5% of which came from the gold mining industry. Of all cases of silicosis, occasional silicotic nodules were found in 44.7% of cases, a few in 30.8%, a moderate number in 17.5% and a large number in 7.0%.

The distribution of cases with silicosis by commodity and population group is presented in Table 4-1.

TABLE 4-1 NUMBER OF CASES AND PREVALENCE OF SILICOSIS BY COMMODITY AND POPULATION GROUP (1986)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	190	98	200	235	0	-	390	139
Platinum	4	22	2	69	0	-	6	28
Coal	5	23	3	38	0	-	8	27
Asbestos	0	-	0	-	2	50	2	23
Iscor	0	-	1	14	0	-	1	13
Diamond	1	59	2	200	0	-	3	111
Copper	0	-	3	176	0	-	3	120
Other	0	-	1	143	0	-	1	67
Unknown	2	43	1	24	0	-	3	34
Total	202	82	213	189	2	44	417	115

Although the silicosis rates increased with increasing age in both black and white men, the age distribution of cases with silicosis differed between the two population groups (Table 4-2).

TABLE 4-2 NUMBER OF CASES AND PREVALENCE OF SILICOSIS IN THE GOLD MINING INDUSTRY, BY AGE AND POPULATION GROUP (1986)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	2	3	0	-	0	-	2	3
30-39	40	68	1	24	0	-	41	66
40-49	86	217	13	165	0	-	99	208
50-59	45	204	33	183	0	-	78	195
60-69	12	273	67	272	0	-	79	272
70-79	1	143	74	347	0	-	75	341
80+	0	-	11	282	0	-	11	282
Unknown	4	82	1	167	0	-	5	91
Total	190	98	200	235	0	-	390	139

Silicosis was diagnosed in men who were young (<40 years) and in men who were exposed to silica for a few years (< 10 years) (Table 4-3).

TABLE 4-3 NUMBER OF CASES AND PREVALENCE OF SILICOSIS IN THE GOLD MINING INDUSTRY, BY YEARS OF SERVICE AND POPULATION GROUP (1986)

Years of service	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
<1	5	31	0	-	0	-	5	30
1-5	20	33	4	73	0	-	24	36
6-10	49	111	4	80	0	-	53	108
11-15	34	214	9	129	0	-	43	188
16-20	24	300	15	208	0	-	39	257
21-25	7	292	26	257	0	-	33	264
26-30	7	350	41	353	0	-	48	353
31-35	0	-	51	375	0	-	51	364
36-40	0	-	35	267	0	-	35	267
41+	1	500	13	224	0	-	14	233
Unknown	43	96	2	35	0	-	45	89
<i>Total</i>	190	98	200	235	0		390	139

SECTION 5 – OTHER PNEUMOCONIOSES

MASSIVE FIBROSIS

There were 12 (0.3%) cases of massive fibrosis (8 black, 4 white). Seven cases of massive fibrosis were from gold, 2 from coal and 3 from the asbestos mining industries.

COAL WORKERS' PNEUMOCONIOSIS

There were 16 (0.4%) cases of coal workers' pneumoconiosis of which 15 cases were from the coal mining industry and 1 case was from the platinum mining industry.

MIXED DUST PNEUMOCONIOSIS

There were 17 (0.5%) cases of mixed dust pneumoconiosis. These cases came from the gold (n=12), coal (n=1) and asbestos (1), platinum (1) and diamond (1) mining industries. One case was from Iscor.

ASBESTOSIS AND PLEURAL PLAQUES

There were 73 cases of asbestosis of which 75.3% (n=55) had slight, 21.9% (n=16) moderate and 2.8% (n=2) marked fibrosis. Of these, 53 (72.6%) had worked in the asbestos mining industry at some time in their lives. There were 21 cases that had asbestos plaques and 12 (57.1%) of these had asbestosis. Note that the parietal pleura (the site where plaque formation usually occurs) is seldom submitted with the lungs.

The distribution of asbestosis by age and population group is shown in Table 5-1.

TABLE 5-1 NUMBER OF CASES AND PREVALENCE OF ASBESTOSIS BY AGE AND POPULATION GROUP (1986)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
30-39	3	4	0	-	0	-	3	4
40-49	9	18	1	8	1	250	11	17
50-59	10	32	4	16	6	857	20	35
60-69	1	14	7	21	13	813	21	50
70-79	1	77	3	12	5	714	9	33
80+	0	-	0	-	4	667	4	74
Unknown	2	29	3	231	0	-	5	60
Total	26	11	18	16	29	644	73	20

SECTION 6 – EMPHYSEMA

There were 977 cases of emphysema, the extent of which was mild in 72.4% (n=707), moderate in 24.9% (n=243) and marked in 2.8% (n=27). The distribution of emphysema by age and population group is presented in Table 6-1.

TABLE 6-1 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY AGE AND POPULATION GROUP (1986)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	17	22	0	-	0	-	17	21
30-39	75	108	4	71	1	333	80	106
40-49	95	185	40	331	0	-	135	211
50-59	80	260	138	552	3	429	221	391
60-69	23	315	237	723	9	563	269	645
70-79	5	385	195	768	6	857	206	752
80+	1	500	31	674	5	833	37	685
Unknown	8	114	4	308	0	-	12	145
Total	304	124	649	577	24	533	977	269

The majority of black and white men with emphysema were from the gold mining industry (72.5%, n=708) (Table 6-2).

TABLE 6-2 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY COMMODITY AND POPULATION GROUP (1986)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	211	108	497	585	0	-	708	253
Platinum	21	115	16	552	0	-	37	175
Coal	51	233	48	615	0	-	99	331
Asbestos	4	174	9	391	23	575	36	419
Iscor	3	429	44	629	0	-	47	610
Diamond	2	118	4	400	0	-	6	222
Copper	0	-	4	235	1	500	5	200
Other	2	250	6	857	0	-	8	533
Unknown	10	217	21	512	0	-	31	352
Total	304	124	649	577	24	533	977	269

TABLE 6-3 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY YEARS OF SERVICE AND POPULATION GROUP (1986)

Years of service	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
<1	3	16	1	200	0	-	4	21
1-5	52	69	20	247	3	429	75	89
6-10	56	104	37	457	6	500	99	157
11-15	29	148	52	565	2	400	83	283
16-20	19	196	65	596	4	667	88	415
21-25	10	278	80	635	2	667	92	558
26-30	11	440	99	678	1	500	111	642
31-35	3	500	102	634	2	1000	107	633
36-40	0	-	106	731	1	1000	107	723
41+	1	500	46	719	0	-	47	701
Unknown	120	196	41	357	3	500	164	224
<i>Total</i>	304	124	649	577	24	533	977	269

SECTION 7 – MESOTHELIOMA

There were 17 cases of mesothelioma in 1986.

TABLE 7-1 NUMBER AND PROPORTION OF MESOTHELIOMA CASES BY AGE AND POPULATION GROUP (1986)

Age group (years)	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
30-39	2	40.0	0	-	0	-	2	11.8
40-49	1	20.0	1	9.1	1	100.0	3	17.6
50-59	2	40.0	0	-	0	-	2	11.8
60-69	0	-	3	27.3	0	-	3	17.6
70-79	0	-	5	45.5	0	-	5	29.4
80-89	0	-	1	9.1	0	-	1	5.9
Unknown	0	-	1	9.1	0	-	1	5.9
<i>Total</i>	5		11		1		17	

The distribution of mesothelioma by commodity and population group is presented in Table 7.2. Four (23.5%) had worked in the asbestos mining industry.

TABLE 7-2 NUMBER AND PROPORTION OF MESOTHELIOMA CASES BY COMMODITY AND POPULATION GROUP (1986)

Commodity	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
Gold	1	20.0	0	-	0	-	9	52.9
Platinum	0	-	0	-	0	-	1	5.9
Coal	1	20.0	0	-	0	-	1	5.9
Asbestos	2	40.0	0	-	1	100.0	3	17.6
Other	0	-	0	-	0	-	1	5.9
Unknown	0	-	2	18.2	0	-	2	11.8
<i>Total</i>	5		11		1		17	

SECTION 8 – PRIMARY LUNG CANCER

Eighty eight cases of primary lung cancer were found at autopsy, 33.0% of which were in black, 61.4% in white and 5.6% in coloured men. Most of the cases were small cell lung carcinomas (39.8%; n=35), followed by squamous lung carcinomas (23.9%; n=21), adeno carcinoma (17.0%; n=15), large cell lung carcinoma (11.4%; n=10), and broncho-alveolar carcinoma (7.9%; n=7)

The distribution of primary lung cancer by age and population group is presented in Table 8-1.

TABLE 8-1 NUMBERS OF CASES AND PREVALENCE OF PRIMARY LUNG CANCER BY AGE AND POPULATION GROUP (1986)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	0	-	1	19	0	-	1	1
30-39	2	3	0	-	0	-	2	3
40-49	10	19	2	17	0	-	12	19
50-59	12	39	11	44	3	429	26	46
60-69	4	55	18	55	2	125	24	58
70-79	1	77	18	71	0	-	19	69
80+	0	-	2	43	0	-	2	37
Unknown	0	-	2	154	0	-	2	24
Total	29	12	54	48	5	111	88	24

The distribution of primary lung cancer by commodity and population group is presented in Table 8-2. The majority of cases came from the gold mining industry.

TABLE 8-2 NUMBER OF CASES AND PREVALENCE OF PRIMARY LUNG CANCER BY COMMODITY AND POPULATION GROUP (1986)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	16	8	33	39	0	-	49	18
Platinum	1	5	1	34	0	-	2	9
Coal	4	18	5	64	0	-	9	30
Asbestos	2	87	2	87	4	100	8	93
Iscor	0	0	6	86	0	-	6	78
Diamond	3	176	0	-	0	-	3	111
Copper	0	-	0	-	1	500	1	40
Other	1	125	2	286	0	-	3	200
Unknown	2	43	5	122	0	-	7	80
Total	29	12	54	48	5	111	88	24

SECTION 9 – CLINICAL CAUSES OF DEATH

Table 9-1 and Figure 9-1 show the clinical causes of death as stated in the accompanying documents submitted with the cardio-respiratory organs, by population group. Diseases of the cardio-vascular system were the most frequent (14.6%) overall. Black men had the highest proportion of unnatural causes of death (66.4%). In 5.0% of all cases, the cause of death was not stated.

TABLE 9-1 CLINICAL CAUSES OF DEATH BY POPULATION GROUP (1986)

System	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
Respiratory	210	8.6	175	15.6	20	44.4	405	11.2
Cardio-vascular	95	3.9	421	37.4	12	26.7	528	14.6
Central Nervous System	86	3.5	62	5.5	2	4.4	150	4.1
Gastro-intestinal	123	5.0	64	5.7	4	8.9	191	5.3
Genito-urinary	34	1.4	27	2.4	0	-	61	1.7
Haematological	18	0.7	10	0.9	0	-	28	0.8
Unnatural	1 632	66.4	223	19.8	4	8.9	1 859	51.3
Miscellaneous	147	6.0	76	6.8	1	2.2	224	6.2
Not stated	111	4.5	67	6.0	2	4.4	180	5.0
Total	2 456		1 125		45		3 626	




FIGURE 9-1 CLINICAL CAUSE OF DEATH AS STATED BY THE CLINICIANS WHO SUBMIT THE ORGANS OF THE DECEASED TO THE NIOH (1986)