



**NATIONAL HEALTH
LABORATORY SERVICE**

NATIONAL INSTITUTE FOR OCCUPATIONAL HEALTH

Pathology Division Surveillance Report

Demographic Data and Disease Rates for January to December 1992

Ntombizodwa Ndlovu

Tony Davies

Gill Nelson

Jill Murray

*PO Box 4788
Johannesburg
2000*

e-mail: jill.murray@nioh.nhls.ac.za

NIOH Report 25/2011

http://www.nioh.ac.za/publications/publications_pathaut_reports.htm

ACKNOWLEDGEMENTS

We thank all the staff members of the NIOH Pathology Department, for their invaluable contribution to the autopsy service.

We also thank Prof Tony Davies for editing the manuscript.

National Institute for Occupational Health, PO Box 4788, Johannesburg, 2000, South Africa

ISSN 1812-7681

EXECUTIVE SUMMARY

During 1992, 3 263 cases came to autopsy at the NIOH. Of these, 65.7% were black men, 32.1% were white and 2.1% were coloured.

Overall disease rates (per 1000 autopsies) for 1992 are shown in Figure 1.




FIG 1 OVERALL DISEASE RATES FOR 1992

TABLE OF CONTENTS

EXECUTIVE SUMMARY	<i>i</i>
GLOSSARY	<i>iv</i>
SECTION 1: BACKGROUND	1
SECTION 2: DEMOGRAPHIC DATA	2
SECTION 3: ACTIVE TUBERCULOSIS	7
SECTION 4: SILICOSIS	9
SECTION 5: OTHER PNEUMOCONIOSES	
<i>Massive Fibrosis</i>	12
<i>Coal Workers' Pneumoconiosis</i>	12
<i>Mixed Dust Pneumoconiosis</i>	12
<i>Asbestosis and pleural plaques</i>	12
SECTION 6: EMPHYSEMA	13
SECTION 7: MESOTHELIOMA	15
SECTION 8: PRIMARY LUNG CANCER	16
SECTION 9: CLINICAL CAUSES OF DEATH	17

LIST OF TABLES

Table 2.1	<i>Distribution of autopsies by year and population group (1992)</i>	2
Table 2.2	<i>Number and proportion of autopsies by type and population group (1992)</i>	3
Table 2.3	<i>Number and proportion of autopsies by age and population group (1992)</i>	3
Table 2.4	<i>Number and proportion of autopsies by commodity and population group (1992)</i>	4
Table 2.5	<i>Number and proportion of autopsies by years of service and population group (1992)</i>	5
Table 2.6	<i>Mean age by commodity and population group (1992)</i>	6
Table 2.7	<i>Mean duration of service by commodity and population group (1992)</i>	7
Table 3.1	<i>Number of cases and prevalence of active PTB by commodity and population group (1992)</i>	9
Table 3.2	<i>Number of cases and prevalence of active PTB by age and population group (1992)</i>	9
Table 4.1	<i>Number of cases and prevalence of silicosis by commodity and population group (1992)</i>	10
Table 4.2	<i>Number of cases and prevalence of silicosis in the gold mining industry, by age and population group (1992)</i>	10
Table 4.3	<i>Number of cases and prevalence of silicosis in the gold mining industry, by years of service and population group (1992)</i>	11
Table 5.1	<i>Number of cases and prevalence of asbestosis by age and population group (1992)</i>	12

Table 6.1	<i>Number of cases and prevalence of emphysema by age and population group (1992)</i>	13
Table 6.2	<i>Number of cases and prevalence of emphysema by commodity and population group (1992)</i>	13
Table 6.3	<i>Number of cases and prevalence of emphysema by years of service and population group (1992)</i>	14
Table 7.1	<i>Number and proportion of mesothelioma cases by age and population group (1992)</i>	15
Table 7.2	<i>Number and proportion of mesothelioma cases by commodity and population group (1992)</i>	15
Table 8.1	<i>Number of cases and prevalence of primary lung cancer by age and population group (1992)</i>	16
Table 8.2	<i>Number of cases and prevalence of primary lung cancer by commodity and population group (1992)</i>	16
Table 9.1	<i>Clinical causes of death by population group (1992)</i>	17

LIST OF FIGURES

Figure 1	<i>Overall disease rates for 1994</i>	<i>i</i>
Figure 2.1	<i>Distribution of autopsies by age and population group (1994)</i>	4
Figure 2.2	<i>Distribution of autopsies by commodity and population group (1994)</i>	5
Figure 2.3	<i>Distribution of autopsies by years of service and population group (1994)</i>	6
Figure 3.1	<i>Distribution of active TB by site (1994)</i>	8
Figure 9.1	<i>Clinical cause of death as given by the clinicians who submit the organs to the NIOH (1994)</i>	17

GLOSSARY

Asbestosis	<i>Lung fibrosis caused by asbestos fibres</i>
Coal workers' pneumoconiosis	<i>Lung fibrosis caused by exposure to coal dust</i>
Emphysema	<i>Lung disease caused by the destruction of the alveolar walls</i>
Massive fibrosis	<i>Lung fibrosis caused by exposure to dust and measuring more than 1 cm in diameter</i>
Mesothelioma	<i>A malignant tumour of the pleural cavity of the lungs</i>
Miner	<i>A person who has worked in a controlled mine or works</i>
Mixed exposures	<i>The multiple dust types to which a miner may be exposed, having worked in several mining commodities in his lifetime</i>
Prevalence	<i>The number of cases in a defined population at a given time</i>
Silicosis	<i>Lung fibrosis caused by inhalation of silica dust; detected by the presence of silicotic nodules in the lung tissue</i>
Surveillance	<i>The ongoing and systematic collection, analysis and interpretation of data related to adverse health outcomes</i>

SECTION 1 – BACKGROUND

The Occupational Diseases in Mines and Works Act, 1973 (Act 78 of 1973) requires that the cardio-respiratory organs of a deceased person who has worked at a controlled mine or a controlled works be examined for the presence of occupational disease, regardless of the clinical cause of death and provided that the next of kin agrees. These examinations are performed by pathologists at the National Institute for Occupational Health (NIOH). A detailed report on each case examined is sent to the Medical Bureau for Occupational Diseases (MBOD). Cases certified as having a compensable disease are then referred to the Compensation Commissioner's office, where the payment for compensation is managed.

Since 1975, the pathological findings from the autopsy examinations have been recorded on the computerised PATHAUT database. PATHAUT comprises data from autopsy examinations and clinical files which include occupational histories. The database is unique and provides an important resource for both surveillance and research. These data are the only comprehensive surveillance data on occupational lung disease in the South African mining industry.

The data presented in this report summarise the PATHAUT system surveillance results, i.e. the results of the systematic collection, collation, and analysis of the pathology findings in the cardio-respiratory organs of mine workers. Data from PATHAUT are exported into, and analysed, using SAS v9.1.

This report describes autopsy cases examined during the year 1992. This and other annual reports can be accessed at www.nioh.ac.za.

Many of the cases had "mixed" exposures in that they had been employed in more than one commodity. For simplicity, cases are categorised according to the commodity in which most years of service were recorded, i.e. the commodity in which the miners had worked for the longest period.

All disease rates reported in this document are expressed per 1000. In all calculations, the denominators used are the total numbers of autopsies in specific commodities, age groups or population groups. Some of these rates must be viewed with caution, as the denominators are very small. This applies, for example, to those commodities where few workers are employed (such as manganese mining), and to the older age groups in some instances.

SECTION 2 – DEMOGRAPHIC DATA

The number of autopsies performed for 1992 is presented in Table 2-1.

TABLE 2-1 DISTRIBUTION OF AUTOPSIES BY YEAR AND POPULATION GROUP (1975-1992)

Year of autopsy	Black		White		Coloured		Unknown		Total N
	N	%	N	%	N	%	N	%	
1975	2 190	71	854	28	32	1			3 076
1976	2 335	68	1 072	31	27	1			3 434
1977	2 351	69	1 039	30	33	1			3 423
1978	2 245	67	1 090	32	32	1			3 367
1979	2 118	66	1 026	33	45	1			3 189
1980	2 338	64	1 274	35	46	1			3 658
1981	2 209	66	1 117	33	33	1			3 359
1982	2 312	63	1 302	36	44	1	1		3 659
1983	2 096	65	1 109	34	41	1			3 246
1984	1 966	64	1 098	36	28	1			3 092
1985	2 275	64	1 200	34	66	2			3 541
1986	2 456	68	1 125	31	45	1			3 626
1987	2 594	68	1 168	30	78	2			3 840
1988	2 518	67	1 165	31	77	2			3 760
1989	2 138	65	1 090	33	60	2			3 288
1990	2 172	64	1 155	34	51	2			3 378
1991	2 143	65	1 080	33	66	2			3 289
1992	2 144	66	1 049	32	70	2			3 263
Total	40 600	66	20 013	33	874	1	1		61 488

It is important to note that a referral bias exists: there is a low autopsy rate for black men who have left employment at the mines, whereas the majority of white retired miners come to autopsy.

The pathologists at the NIOH perform two types of autopsy examinations. For men dying distant from Johannesburg, the cardio-respiratory organs are removed locally, preserved in formalin and sent to the NIOH. Full autopsies are undertaken on men who die close to Johannesburg.

Table 2-2 shows the distribution of autopsies by population group for 1992. Autopsies of only the cardio-respiratory organs comprised 89.2% of all examinations.

TABLE 2-2 NUMBER AND PROPORTION OF AUTOPSIES BY TYPE AND POPULATION GROUP (1992)

Autopsy type	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
<i>Cardiorespiratory organs only</i>	2 057	95.9	783	74.6	69	98.6	2 909	89.2
<i>Full autopsy</i>	83	3.9	264	25.2	0	0.0	347	10.6
<i>Not stated</i>	4	0.2	2	0.2	1	1.4	7	0.2
<i>Total</i>	2 144		1 049		70		3 263	

The age distributions of autopsies for 1992 are shown in Table 2-3 and Figure 2-1. The mean age at autopsy of black men was 38.0 years. The mean age of white men at autopsy was 60.3 years and for coloured men 64.1 years

TABLE 2-3 NUMBER AND PROPORTION OF AUTOPSIES BY AGE AND POPULATION GROUP (1992)

Age group (years)	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
<20	1	-	3	0.3	0	-	4	0.1
20-29	400	18.7	39	3.7	1	1.4	440	13.5
30-39	678	31.6	83	7.9	0	-	761	23.3
40-49	557	26.0	104	9.9	11	15.7	672	20.6
50-59	331	15.4	191	18.2	16	22.9	538	16.5
60-69	77	3.6	289	27.6	17	24.3	383	11.7
70-79	7	0.3	256	24.4	13	18.6	276	8.5
80+	2	0.1	78	7.4	12	17.1	92	2.8
<i>Unknown</i>	91	4.2	6	0.6	0	-	97	3.0
<i>Total</i>	2 144		1 049		70		3 263	




FIG 2-1 DISTRIBUTION OF AUTOPSIES BY AGE AND POPULATION GROUP (1992)

Cases were placed in categories according to the commodity in which they had worked for the longest duration (most exposure). Many men worked in a number of different mining commodities during their lifetimes and had “mixed” exposures. This was not taken into account in the analysis of exposure type (commodity).

Table 2-4 and Figure 2-2 show the distribution of autopsies by commodity and population group for 1992.

TABLE 2-4 NUMBER AND PROPORTION OF AUTOPSIES BY COMMODITY AND POPULATION GROUP (1992)

Commodity	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
Gold	1 662	77.5	776	74.0	0	-	2 438	74.7
Platinum	259	12.1	48	4.6	0	-	307	9.4
Coal	123	5.7	71	6.8	1	1.4	195	6.0
Asbestos	18	0.8	39	3.7	61	87.1	118	3.6
Iscor	4	0.2	41	3.9	0	-	45	1.4
Diamond	29	1.4	16	1.5	1	1.4	46	1.4
Copper	5	0.2	21	2.0	6	8.6	32	1.0
Other	2	0.1	4	0.4	0	-	6	0.2
Unknown	42	2.0	33	3.1	1	1.4	76	2.3
Total	2 144		1 049		70		3 263	

NOTE: This table shows only those commodities where a total of 10 or more cases were received





FIG 2-2 DISTRIBUTION OF AUTOPSIES BY COMMODITY AND POPULATION GROUP (1992)

Detailed information about the years in mining service by population group is presented in Table 2-5 and Figure 2-3.

TABLE 2-5 NUMBER AND PROPORTION OF AUTOPSIES BY YEARS OF SERVICE AND POPULATION GROUP (1992)

Years of service	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
<1	83	3.9	5	0.5	1	1.4	89	2.7
1-5	408	19.0	79	7.5	11	15.7	498	15.3
6-10	413	19.3	95	9.1	23	32.9	531	16.3
11-15	406	18.9	122	11.6	17	24.3	545	16.7
16-20	176	8.2	144	13.7	8	11.4	328	10.1
21-25	60	2.8	134	12.8	2	2.9	196	6.0
26-30	26	1.2	128	12.2	3	4.3	157	4.8
31-35	11	0.5	123	11.7	1	1.4	135	4.1
36-40	2	0.1	67	6.4	0	-	69	2.1
41+	3	0.1	36	3.4	0	-	39	1.2
Unknown	556	25.9	116	11.1	4	5.7	676	20.7
Total	2 144		1 049		70		3 263	



Years of service:

■ < 1
 ■ 1 - 5
 ■ 6 - 10
 ■ 11 - 15
 ■ 16 - 20
 ■ 21 - 25
 ■ 26 - 30
 ■ 31 - 35
 ■ 36 - 40
 ■ 41+

FIG 2-3 DISTRIBUTION OF AUTOPSIES BY YEARS OF SERVICE AND POPULATION GROUP (1992)

The mean age and duration of service by commodity type and population group for those cases for which information was available are shown in Tables 2-6 and 2-7.

TABLE 2-6 MEAN AGE BY COMMODITY AND POPULATION GROUP (1992)

Commodity	Black			White		
	N	Mean (years)	SD*	N	Mean (years)	SD*
Gold	1 589	39.7	10.4	774	61.4	15.3
Platinum	254	37.1	9.9	48	54.1	14.8
Coal	113	41.5	12.2	71	62.1	15.2
Asbestos	18	48.6	12.1	39	60.6	11.1
Iscor	4	50.3	13.8	41	56.3	15.5
Diamond	29	41.6	12.0	15	51.4	13.9
Copper	5	55	7.8	21	61.0	16.5
Other	2	48.5	2.1	4	59.3	6.2
Unknown	39	43.5	11.9	31	58.1	16.4
Total	2 053	39.7	10.7	1 044	60.6	15.3

* Standard deviation

TABLE 2-7 MEAN DURATION OF SERVICE BY COMMODITY AND POPULATION GROUP (1992)

Commodity	Black			White		
	N	Mean (years)	SD*	N	Mean (years)	SD*
<i>Gold</i>	1 307	10.4	6.9	708	23.2	11.2
<i>Platinum</i>	173	8.0	4.6	43	16.5	8.4
<i>Coal</i>	60	10.2	5.9	63	19.5	19.5
<i>Asbestos</i>	9	10.9	6.4	32	13.6	9.3
<i>Iscor</i>	3	4.4	2.8	32	18.6	11.9
<i>Diamond</i>	21	12.7	9.8	14	15.8	6.7
<i>Copper</i>	0	-	-	18	19.0	9.7
<i>Other</i>	2	19.3	3.8	4	23.5	8.2
<i>Unknown</i>	13	13.2	11.2	19	15.4	10.4
<i>Total</i>	1 588	10.2	6.7	933	21.8	11.2

* Standard deviation

SECTION 3 – ACTIVE TUBERCULOSIS

The distribution of active tuberculosis (TB) by anatomical site is presented in Figure 3-1 (n=226). Active pulmonary TB (PTB) was diagnosed in 5.2% (n=169) of all cases autopsied in 1992. Most of the men with PTB were black (89.3%; 151 cases), 7.1% (12 cases) were white and 3.6% (6 cases) were coloured.




FIG 3-1 DISTRIBUTION OF ACTIVE TB BY SITE (1992)

The distribution of active PTB cases by commodity is shown in Table 3-1. The rates in this and subsequent tables and figures are expressed per 1000. The majority of active PTB cases (86.4%) came from the gold mining industry (74.7% of all autopsy cases came from that commodity).

TABLE 3-1 NUMBER OF CASES AND PREVALENCE OF ACTIVE PTB BY COMMODITY AND POPULATION GROUP (1992)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	135	81	11	14	0	-	146	60
Platinum	9	35	0	-	0	-	9	29
Coal	4	33	0	-	0	-	4	21
Asbestos	1	56	1	26	5	82	7	59
Diamond	1	34	0	-	0	-	1	22
Copper	0	-	0	-	1	167	1	31
Unknown	1	24	0	-	0	-	1	13
Total	151	70	12	11	6	86	169	52

The age distribution of cases with active PTB is shown in Table 3-2. Most of the cases were aged between 20 and 69 years (156 cases=92.3%).

TABLE 3-2 NUMBER OF CASES AND PREVALENCE OF ACTIVE PTB BY AGE AND POPULATION GROUP (1992)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	13	33	0	-	0	-	13	30
30-39	41	60	0	-	0	-	41	54
40-49	56	101	1	10	1	91	58	86
50-59	26	79	4	21	1	63	31	58
60-69	8	104	2	7	3	176	13	34
70-79	1	143	4	16	1	77	6	22
80+	0	-	1	13	0	-	1	11
Unknown	6	66	0	-	0	-	6	62
Total	151	70	12	11	6	86	169	52

SECTION 4 – SILICOSIS

Silicotic nodules were found in the lungs of 393 cases (12.0% of all autopsies), 95.7% of which came from the gold mining industry. Of all cases of silicosis, occasional silicotic nodules were found in 45.3% of cases, a few in 24.2%, a moderate number in 20.1%, a large number in 7.4% and in 3.1% of the cases severity was not stated.

The distribution of cases with silicosis by commodity and population group is presented in Table 4-1.

TABLE 4-1 NUMBER OF CASES AND PREVALENCE OF SILICOSIS BY COMMODITY AND POPULATION GROUP (1992)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	224	135	152	196	0	-	376	154
Platinum	3	12	1	21	0	-	4	13
Coal	1	8	4	56	0	-	5	26
Asbestos	0	-	1	26	3	49	4	34
Copper	0	-	1	48	0	-	1	31
Unknown	2	48	1	30	0	-	3	39
Total	230	107	160	153	3	43	393	120

Although the silicosis rates increased with increasing age in both black and white men, the age distribution of cases with silicosis differed between the two population groups (Table 4-2).

TABLE 4-2 NUMBER OF CASES AND PREVALENCE OF SILICOSIS IN THE GOLD MINING INDUSTRY, BY AGE AND POPULATION GROUP (1992)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	5	17	0	-	0	-	5	15
30-39	29	55	2	38	0	-	31	53
40-49	90	200	5	71	0	-	95	183
50-59	80	310	15	109	0	-	95	241
60-69	13	277	41	194	0	-	54	209
70-79	1	200	65	311	0	-	66	308
80+	0	-	23	377	0	-	23	371
Unknown	6	82	1	500	0	-	7	93
Total	224	135	152	196	0	-	376	154

Silicosis was diagnosed in men who were young (<40 years) and in men who were exposed to silica for a few years (< 10 years) (Table 4-3).

TABLE 4-3 NUMBER OF CASES AND PREVALENCE OF SILICOSIS IN THE GOLD MINING INDUSTRY, BY YEARS OF SERVICE AND POPULATION GROUP (1992)

Years of service	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
<1	8	110	0	-	0	-	8	105
1-5	17	53	0	-	0	-	17	46
6-10	25	76	6	98	0	-	31	79
11-15	62	183	14	175	0	-	76	182
16-20	50	323	23	223	0	-	73	283
21-25	13	232	28	269	0	-	41	256
26-30	5	238	23	221	0	-	28	224
31-35	3	375	37	346	0	-	40	348
36-40	2	1000	13	217	0	-	15	242
41+	1	333	6	176	0	-	7	189
Unknown	38	107	2	29	0	-	40	95
<i>Total</i>	224	135	152	196	0		376	154

SECTION 5 – OTHER PNEUMOCONIOSES

MASSIVE FIBROSIS

There were 21(0.6%) cases of massive fibrosis (15 black and 6 white). Seventeen cases of massive fibrosis were from gold, two from platinum and for two cases the commodity was not known.

COAL WORKERS' PNEUMOCONIOSIS

There were 21 (0.6%) cases of coal workers' pneumoconiosis of which 15 cases were from the coal, four from gold, and one each from the platinum and asbestos mining industries. For one case the commodity was not known.

MIXED DUST PNEUMOCONIOSIS

There were 11 (0.3%) cases of mixed dust pneumoconiosis. These cases came from the gold (n=10) and coal (n=1) mining industries.

ASBESTOSIS AND PLEURAL PLAQUES

There were 100 cases of asbestosis of which 63.0% (n=63) had slight, 35% (n=35) moderate and 2% (n=2) marked fibrosis. Of these, 73 (73.0%) had worked in the asbestos mining industry at some time in their lives. There were 27 cases that had asbestos plaques and 16 (59.3%) of these had asbestosis. Note that the parietal pleura (the site where plaque formation usually occurs) is seldom submitted with the lungs.

The distribution of asbestosis by age and population group is shown in Table 5-1.

TABLE 5-1 NUMBER OF CASES AND PREVALENCE OF ASBESTOSIS BY AGE AND POPULATION GROUP (1992)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	1	3	1	26	0	-	2	5
30-39	3	4	0	-	0	-	3	4
40-49	8	14	2	19	4	364	14	21
50-59	16	48	6	31	8	500	30	56
60-69	5	65	10	35	11	647	26	68
70-79	0	-	9	35	7	538	16	58
80+	0	-	2	26	6	500	8	87
Unknown	1	11	0	-	0	-	1	10
Total	34	16	30	29	36	514	100	31

SECTION 6 – EMPHYSEMA

There were 664 cases of emphysema, the extent of which was mild in 68.2% (n=453), moderate in 27.0% (n=179) and marked in 4.8% (n=32). The distribution of emphysema by age and population group is presented in Table 6-1.

TABLE 6-1 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY AGE AND POPULATION GROUP (1992)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
20-29	3	8	0	-	0	-	3	7
30-39	41	60	5	60	0	-	46	60
40-49	68	122	12	115	2	182	82	122
50-59	57	172	78	408	4	250	139	258
60-69	21	273	136	471	8	471	165	431
70-79	3	429	160	625	7	538	170	616
80+	0	-	39	500	6	500	45	489
Unknown	12	132	2	333	0	-	14	144
Total	205	96	432	412	27	386	664	203

The majority of black and white men with emphysema were from the gold mining industry (71.7%, n=476) (Table 6-2).

TABLE 6-2 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY COMMODITY AND POPULATION GROUP (1992)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	154	93	322	415	0	-	476	195
Platinum	17	66	12	250	0	-	29	94
Coal	22	179	34	479	0	-	56	287
Asbestos	2	111	19	487	26	426	47	398
Isacor	0	-	16	390	0	-	16	356
Diamond	4	138	5	313	0	-	9	196
Copper	1	200	10	476	1	167	12	375
Other	0	-	2	500	0	-	2	333
Unknown	5	119	12	364	0	-	17	224
Total	205	96	432	412	27	386	664	203

TABLE 6-3 NUMBER OF CASES AND PREVALENCE OF EMPHYSEMA BY YEARS OF SERVICE AND POPULATION GROUP (1992)

Years of service	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
<1	3	36	0	-	0	-	3	34
1-5	20	49	15	190	3	273	38	76
6-10	42	102	26	274	12	522	80	151
11-15	41	101	59	484	7	412	107	196
16-20	20	114	62	431	1	125	83	253
21-25	14	233	69	515	1	500	84	429
26-30	7	269	55	430	2	667	64	408
31-35	3	273	61	496	0	-	64	474
36-40	0	-	39	582	0	-	39	565
41+	0	-	16	444	0	-	16	410
Unknown	55	99	30	259	1	250	86	127
<i>Total</i>	205	96	432	412	27	386	664	203

SECTION 7 – MESOTHELIOMA

There were 13 cases of mesothelioma in 1992.

TABLE 7-1 NUMBER AND PROPORTION OF MESOTHELIOMA CASES BY AGE AND POPULATION GROUP (1992)

Age group (years)	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
30-39	3	50.0	0	-	0	-	3	23.1
40-49	0	-	1	16.7	0	-	1	7.7
50-59	2	33.3	1	16.7	0	-	3	23.1
60-69	1	16.7	1	16.7	0	-	2	15.4
70-79	0	-	2	33.3	1	100.0	3	23.1
Unknown	0	-	1	16.7	0	-	1	7.7
<i>Total</i>	6		6		1		13	

The distribution of mesothelioma by commodity and population group is presented in Table 7.2. Ten (76.9%) had worked in the asbestos mining industry.

TABLE 7-2 NUMBER AND PROPORTION OF MESOTHELIOMA CASES BY COMMODITY AND POPULATION GROUP (1992)

Commodity	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
Gold	1	17.7	3	50.0	0	-	4	30.8
Coal	0	-	1	16.7	0	-	1	7.7
Asbestos	5	83.3	2	33.3	1	100.0	8	61.5
<i>Total</i>	6		6		1		13	

SECTION 8 – PRIMARY LUNG CANCER

Ninety nine cases of primary lung cancer were found at autopsy, 25.3% of which were in black, 63.6% in white and 11.1% in coloured men. Most of the cases were squamous lung carcinomas (41.4%; n=41), followed by large cell lung carcinoma (21.2%; n=21) and small cell lung carcinomas (21.2%; n=21), adeno carcinoma (12.1%; n=12) and broncho-alveolar carcinoma (4.0%; n=4).

The distribution of primary lung cancer by age and population group is presented in Table 8-1.

TABLE 8-1 NUMBERS OF CASES AND PREVALENCE OF PRIMARY LUNG CANCER BY AGE AND POPULATION GROUP (1992)

Age group (years)	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
30-39	6	9	0	-	0	-	6	8
40-49	9	16	1	10	1	91	11	16
50-59	6	18	9	47	4	250	19	35
60-69	3	39	23	80	2	118	28	73
70-79	0	-	23	90	1	77	24	87
80+	0	-	6	77	3	250	9	98
Unknown	1	11	1	167	0	-	2	21
Total	25	12	63	60	11	157	99	30

The distribution of primary lung cancer by commodity and population group is presented in Table 8-2. The majority of cases came from the gold mining industry.

TABLE 8-2 NUMBER OF CASES AND PREVALENCE OF PRIMARY LUNG CANCER BY COMMODITY AND POPULATION GROUP (1992)

Commodity	Black		White		Coloured		Total	
	N	Rate	N	Rate	N	Rate	N	Rate
Gold	16	10	45	58	0	-	61	25
Platinum	5	19	3	63	0	-	8	26
Coal	1	8	4	56	0	-	5	26
Asbestos	1	56	5	128	9	148	15	127
Iscor	0	-	1	24	0	-	1	22
Diamond	1	34	0	-	0	-	1	22
Copper	0	-	0	-	1	167	1	31
Other	0	-	1	250	0	-	1	167
Unknown	1	24	4	121	0	-	5	66
Total	25	12	63	60	10	143	98	30

SECTION 9 – CLINICAL CAUSES OF DEATH

Table 9-1 and Figure 9-1 show the clinical causes of death as stated in the accompanying documents submitted with the cardio-respiratory organs, by population group. Diseases of the cardio-vascular system were the most frequent (10.0%) overall. Black men had the highest proportion of unnatural causes of death (46.9%). In 25.3% of all cases, the cause of death was not stated.

TABLE 9-1 CLINICAL CAUSES OF DEATH BY POPULATION GROUP (1992)

System	Black		White		Coloured		Total	
	N	%	N	%	N	%	N	%
Respiratory	178	8.3	134	12.8	14	20.0	326	10.0
Cardio-vascular	72	3.4	219	20.9	13	18.6	304	9.3
Central Nervous System	71	3.3	37	3.5	4	5.7	112	3.4
Gastro-intestinal	103	4.8	47	4.5	6	8.6	156	4.8
Genito-urinary	35	1.6	25	2.4	1	1.4	61	1.9
Haematological	17	0.8	14	1.3	0	-	31	1.0
Unnatural	1 005	46.9	154	14.7	6	8.6	1 165	35.7
Miscellaneous	169	7.9	102	9.7	11	15.7	282	8.6
Not stated	494	23.0	317	30.2	15	21.4	826	25.3
Total	2 144		1 049		70		3 263	




FIGURE 9-1 CLINICAL CAUSE OF DEATH AS STATED BY THE CLINICIANS WHO SUBMIT THE ORGANS OF THE DECEASED TO THE NIOH (1992)